

Potential of Fruits to Improve Dyslipidemias: A Pilot Review

Miguel Angel Flores Flores^{1,*}, Luis Antonio Flores Flores^{2,*}, Serafín Dernali Romero Yupanqui^{3,*}

¹Faculty of Food Industry, Universidad Nacional de la Amazonía Peruana, Iquitos, Maynas, Loreto, 16002, Perú; ²Faculty of Chemical Engineering, Universidad Nacional de la Amazonía Peruana, Iquitos, Maynas, Loreto, 16002, Perú; ³Faculty of Medicine, Universidad Nacional de la Amazonía Peruana, Iquitos, Maynas, Loreto, 16002, Perú

*These authors contributed equally to this work

Correspondence: Miguel Angel Flores Flores, Faculty of Food Industry, Universidad Nacional de la Amazonia Peruana, Iquitos, Maynas, Loreto, 16002, Perú, Tel +51987282346, Email mianfloflo@gmail.com

Abstract: Dyslipidemia is a condition characterized by excessive lipids in the blood plasma, including triglycerides (TG), total cholesterol (TC), low-density lipoprotein cholesterol (LDL-C) and decreased levels of high-density lipoprotein cholesterol (HDL-C), which is generated mostly due to obesity. This study aims to summarize research conducted on rats and humans regarding the potential of eight fruits in reducing dyslipidemia and their associated health benefits (eg, reduction of free radicals, hypoglycemic effects, weight reduction, lowering of blood pressure, and anti-inflammatory properties). The studied fruits include pomegranate, star fruit, Rosa roxburghii, pineapple, tree tomato, coffee, apple, and passion fruit. Various parts of these fruits, such as the root, leaves, stem, peel, and pulp, were analyzed for their effects. These fruits are edible, widely available, and cost-effective when purchased during the harvest season (Graphical abstract).

Keywords: fruits, dyslipidemias, humans, rats, benefits

Introduction


Overweight and obesity result from an imbalance between excessive energy intake (primarily from foods rich in fat and refined sugars) and insufficient energy expenditure. This leads to excess adipose and visceral tissue accumulation, which can contribute to non-communicable diseases and dyslipidemia.¹ Dyslipidemia is characterized by elevated cholesterol and triglyceride levels, resulting in abnormal lipoprotein levels such as LDL (low-density lipoprotein) and HDL (high-density lipoprotein). The most common consequences associated with dyslipidemia include an increased risk of atherosclerotic cardiovascular disease (ASCVD) and pancreatitis, particularly when triglyceride levels are high.² The treatment of dyslipidemia can be pharmacological or involve natural products, which have gained increasing scientific attention. These natural products include nutritional supplements, fresh fruits, and fruit juices that help inhibit lipid oxidation due to the powerful antioxidant effects of phenolic compounds, oils, and dietary fiber.

This study aims to highlight the bioactive properties of eight fruits (Figure 1) that help prevent dyslipidemia. Their primary nutrients, functional health effects, and mechanisms of action will be discussed. The findings aim to encourage the consumption of fresh fruits as a preventive measure against dyslipidemia and related health issues.

Methods

The information presented in this review was obtained from reputable scientific databases, including ScienceDirect, PubMed, Scopus, and Google Scholar. The keywords used for the literature search included “Apple” or “Malus domestica”, “Pineapple” or “Ananas comosus”, “Tree tomato”, “Solanum betaceum” or “Cythomandra betacea”, “Coffee”, “Pomegranate” or “Punica granatum”, “Starfruit” or “Averrhoa carambola L.”, “Rosa roxburghii” and “Passion fruit” or “Passiflora edulis” in combination

Graphical Abstract



with the words “Dyslipidemia”, “Hyperlipidemia”, “rats” and “Human”. In short, all the information to present in the following review.

Fruits That Prevent the Development of Dyslipidemia in Experimental Models

Pomegranate (*Punica granatum* L.)

Pomegranate is an ancient fruit from the *Punicaceae* family, often referred to as a “superfruit or miracle fruit” due to its high nutritional value, including carbohydrates, vitamins, and minerals.³ It has been cultivated worldwide (since the northern hemisphere to the middle east)⁴ and is particularly known for its cardiovascular benefits, especially in reducing excess cholesterol in the blood.³

Research in rats indicates that ad libitum consumption of fermented pomegranate juice for a period of 6 weeks reduces at least 48% of blood TG.⁴ In another study, they found that the consumption of 250 mg/kg of weight per day for 35 days of pomegranate peel and pulp reduced TC, TG, and LDL levels and increased HDL levels.⁵ On the other hand, consumption of 1 mL of fresh juice significantly reduced plasma CT and TG values, and 100 mg of seed powder consumption raised HDL, all of this was analyzed for 22 days.⁶

Studies in humans with 66 obese women who supplemented with 500 mg of pomegranate peel extract for 8 weeks found a reduction in TC and TG.⁷ Likewise, in another study with 14 volunteers who consumed 300 mL/day of fresh pomegranate juice for 2 weeks, lower LDL levels were evident.⁸ The use of 1 g/day for 2 weeks of pomegranate extract pill (POMx) in 12




Figure 1 Fruits that prevent dyslipidemia. Created in BioRender. Flores Flores, M. (2025) <https://BioRender.com/i31f701>.

volunteers decreased LDL by 26%, reactive oxygen species (ROS) by up to 30%, triglycerides by up to 48%, and cholesterol by up to 44%.⁹ Eleven volunteers with acute coronary syndrome used 20 g/day of microencapsulated pomegranate dissolved in 250 mL of water for 30 days, decreased triglycerides, cholesterol and LDL-c by 42%, 15% and 35% after 8 hours, respectively.¹⁰ Twenty-two women with polycystic ovary syndrome received 45 mL/day combined with 180 mL of water, decreased triglycerides and LDL-c.¹¹ Sixty-six patients on hemodialysis received 100 mL of pomegranate juice 3 times a week for 1 year, and the results showed significantly higher levels of HDL and lower levels of triglycerides.¹² Thirty patients with Type 2 diabetes mellitus (T2DM) received capsules containing 250 mg twice daily per 8 weeks of potential of pomegranate peel extract (PoPEx) twice daily, the results were significant decrease in the levels of TC, LDL-C and TG and significant increase of HDL-C.¹³ 51 hyperlipidemia subjects received 400 mg pomegranate seed oil (PSO) twice daily per 4 weeks, the result was non-significant increase in HDL-C concentration.¹⁴ Twenty-two type II diabetic patients with hyperlipidemia consumed 40 g/day of concentrated pomegranate juice (CPJ) for 8 weeks, the results showed significant reductions of total cholesterol and low-density lipoprotein (LDL)-cholesterol (Figure 2).^{15,16}

This reduction in dyslipidemia is thanks to the bioactive compounds contained in different parts of the pomegranate, such as³ pomegranate juice is rich in polyphenols such as ellagitannins, anthocyanins, flavonoids, and helps modulate the intestinal microbiota with the increased probiotics,⁴ and the seeds rich in conjugated fatty acids and the peel rich in ellagitannins^{17,18} and the peel extract are rich in anthocyanins and tannins,⁷ and the fresh juice is rich in malic acid, polyphenol and tryptophan.^{6,8}

These investigations also indicated that the consumption of pomegranate in its different presentations indicated above also has effects on the reduction of body fat storage, weight reduction,⁴ fatty liver, visceral and subcutaneous,⁵ anti-inflammatory,⁶ reduction of free radicals,⁷ levels of docosahexaenoic acid (22:6n-3, DHA) in erythrocytes, slowed the development of atherosclerotic plaque in the carotid arteries and optimized stress-induced myocardial ischemia in patients with coronary artery disease.⁹

Starfruit (*Averrhoa carambola*)

Star fruit, family of *Oxalidaceae*, is believed to be native to Malaysia and is usually called “sweet belimbing or star fruit” being a natural source of antioxidants and vitamin C that help strengthen immunity and eliminate toxins that affect the body.¹⁹




Figure 2 Bioactive compounds present in Pomegranate (*Punica granatum* L.) and their effect in vivo. Created in BioRender. Flores Flores, M. (2025) <https://BioRender.com/e75a587>.

Notes: Polyphenolic compounds in pomegranate juice increase the sensitivity of insulin receptors, activate γ -PPAR receptors, modulate GLUT-4 expression and thus improve glucose uptake by tissues. Pomegranate extracts can reduce glucose absorption by inhibiting enzymes such as α -amylase and pancreatic lipase.⁴

Aladeileh and Saghir^{19,20} investigated the use of Star fruit leaf extract in rats, reaching the conclusion that regardless of the doses and for a period of 5 weeks, the values of TC, TG, LDL and increase of HDL. 500 mg/kg for 48 hours of ethyl acetate fractions in Star fruit leaves had a greater effect regarding CT and TG.²¹ 300 mg/kg of aqueous methanol extract of the leaves for a period of 4 weeks reduced TC, TG and LDL.²² The use of 120 mg/kg for 21 days of benzoquinone extracted from Star fruit roots reduced CT and TG levels.²³ The ad libitum use of star fruit bagasse for 30 days reduced the concentrations of TG, cholesterol and LDL in mice.²⁴ Also, the polyphenols from the fruit that was applied intragastrically for 14 weeks reduced the levels of CT and TG (Figure 3).²⁵

The extract of Star fruit leaves is rich in ascorbic acid and phenolic compounds such as gallic acid and epicatechin,²⁶ while stem extracts have antioxidant effects thanks to their flavonoid content.^{13,20} Star fruit bagasse is rich in insoluble fiber.²⁴




Figure 3 Bioactive compounds present in Starfruit (*Averrhoa carambola*) and their effect in vivo. BioRender. Flores Flores, M. (2025) <https://BioRender.com/e75a587>.

Notes: In cholesterol synthesis, HMG-CoA reductase reduces HMG-CoA to mevalonate, which is a step in cholesterol synthesis. This is where the bioactive compounds in carambola leaves inhibit the action. Apigenin stimulates the immune system, reducing plasma levels of LDL-C (which were not used) by inhibiting LDL receptors in hepatocytes and expelling cholesterol through the feces.²¹

Abbreviation: HMG-CoA, 3-hydroxy-3-methylglutaryl-coenzyme A.

This reduction in dyslipidemia is associated with the inhibitory effect of HMG-CoA (3-hydroxy-3-methylglutaryl-coenzyme A) reductase in the blood, the activation of pancreatic lipase and the antioxidant effect of GSH (L-g-glutamyl-L-cysteinyl-glycine), the enzyme superoxide dismutase (SOD) and catalase.^{19,21,22}

Research affirms that the consumption of star fruit juice also has anti-inflammatory and antiulcerogenic effects, reduces body weight¹⁹ and helps with diarrhea, vomiting, and hepatic colic.²⁷ While the leaves may have anti-obesity effect²² and fruit stem have antioxidant effect.²⁰ On the other hand, roots are associated with the reduction of inflammatory factors because it has against tumor necrosis factor- α (TNF- α) and interleukin-6 (IL-6).²³ The insoluble fiber found in bagasse is also associated with lowering blood glucose.^{22,24}

Rosa roxburghii (*Rosa roxburghii* Tratt)

The fruit of the *Rosa roxburghii*, family Rosaceae, grows mostly in mountains between 200 and 2500 meters above sea level and is cultivated in southwest China. Fruit rich in vitamin C, which is why it is known as the “king of vitamin C”.²⁸

Studies with 48 rats with dyslipidemia indicate that the use of 10%, 30% and 50% (0.1 mL/10 g of weight) of rose roxburghii fruit vinegar for 30 days decreases the levels of TC, TG and LDL and increases the HDL.²⁸ Likewise, the use of 125, 250 and 500 mg/kg of seed oil for 8 weeks has effects on the inhibition of the accumulation of lipids such as CT and TG.²⁹ 900 mg/kg/day for 8 weeks of polysaccharides isolated from fruit showed a significant decrease in TC, TG, LDL and an increase in HDL.³⁰ Consumption of 10 mL/kg/day for 12 weeks of fermented juice decreases the levels of TC, TG, LDL and increases HDL which are associated with beneficial intestinal bacteria.³¹ 120 mg/kg of hydroalcoholic extract of the fruit for 8 weeks helped to decrease TC, TG, LDL and increase HDL.³²

Jin et al³³ studied 100 patients with psychiatric disorders and antipsychotic-induced metabolic syndrome (MetS), of which, 50 received daily dose of 32 mg of Chiglitazar combined with *Rosa roxburghii* Tratt per 12 weeks; the results were significant decreases in triglyceride (TG) and total cholesterol (TC) levels (Figure 4).

The fermented juice and fresh juice are rich in flavonoids, polyphenols, and polysaccharides in general and rich in vitamin C.³¹ Fruit vinegar is rich in acetic acid, flavonoids and total phenolics,²⁸ while seed oil is rich in unsaturated fatty acids, including linoleic acid, α -linolenic acid and oleic acid.²⁹ Likewise, the hydroalcoholic extract is rich in phenolic acid³² and the fruit itself is rich in polysaccharides such as organic acid, triterpenes and superoxide dismutase.³⁰

In vivo research indicates that the polysaccharide of rose roxburghii fruit has an antioxidant effect, which acts by reducing lipid peroxidation (LPO) and malondialdehyde (MDA), which is used as a biomarker of oxidative stress,³⁴ inhibiting the activity of α -D-glucosidase, decreasing the formation of glucose, which prevents postprandial blood sugar levels from rising in patients with diabetes.³⁵ Polysaccharides and fermented juice reduce colon bacteria such as *Firmicutes/Bacteroidetes* responsible for the risk of diabetes and dyslipidemia.^{30,31} The vinegar from the rose roxburghii fruit influences weight reduction, helps improve hypertension and diabetes, and regulates intestinal flora by increasing *Akkermansia* and *Bacteroides* bacteria.²⁸ The seed oil also helps in increasing gene expression of fatty acid oxidation, biogenesis and mitochondrial function for the prevention of non-alcoholic fatty liver.²⁹ The hydroalcoholic extract also helps inhibit lipoprotein lipase and the activity of antioxidant enzymes.³² Polyphenols of fresh juice exhibit protective effects against lung injury by decreasing inflammatory cytokine levels and affecting key metabolic pathways, highlighting their potential therapeutic role in mitigating inflammation-related complications.³³

Pineapple (*Ananas comosus*)

Pineapple is a fruit that belongs to the *Bromeliaceae* family and is mostly cultivated in equatorial areas due to its medicinal properties on the health of people suffering from tumors, helps wound healing³⁶ and aids in digestion.³⁷

Studies in rats indicate that consuming 200 mg/kg/day for 8 weeks of dehydrated pineapple pulp and fresh fruit reduced TC, TG and LDL levels.^{38,39} A 15% dose of pineapple juice for 8 weeks significantly reduced TG, TC, LDL and VLDL levels and increased HDL.⁴⁰ 0.40 mg/kg of ethanolic extract of pineapple leaves for 15 days inhibits the increase in postprandial TG, TC and LDL.⁴¹ Treatment with 20 mg/kg day of bromelain from fruit, leaves, and fruit peel for 4 weeks reduced lipid accumulation and therefore the reduction of CT and TG (Figure 5).⁴²

The pineapple stem is rich in bromelain and has anticancer, anti-inflammatory, antiplatelet, antioxidant and significant weight reduction effects.^{36,42} The pulp is rich in phenols, flavonoids, gallic acid, catechin, epicatechin and ferulic acid, so its contribution to health is also as an antioxidant and helps in weight reduction;³⁸ pineapple juice, rich in phenolic compounds such as: gallic acid, gentisic acid, isoferulic acid, vanillin, ferulic acid, sinapic acid, syringic acid and o-coumaric acid that help reduce visceral fat.⁴⁰ Phenols in pineapple leaves have effects on reducing abdominal fat, hypoglycemic and reducing fatty liver.³⁷

Tree Tomato (*Solanum betaceum* / *Cyphomandra betacea*)

The tree tomato, also known as “tomatillo”, “tamarillo” “Chilto”, belonging to the *Solanaceae* family, is produced worldwide, as well as in Latin America. It is a highly valued fruit due to its high vitamin content and among them the phytochemicals that include anthocyanins, flavonoids, phenolic acids, ascorbic acid⁴³ and fiber.⁴⁴

Studies on rats with a dose of 150, 200 and 300 mg/kg of tree tomato powder extract for 7 weeks significantly reduced the level of TG, LDL and increased HDL.⁴³ In another study, they used 100, 200 and 400 mg/kg/day of dried




Figure 4 Bioactive compounds present in *Rosa roxburghii* (*Rosa roxburghii tratt*) and their effect in vivo. Created in BioRender. Flores Flores, M. (2025) <https://BioRender.com/171gl60>.

Notes: Dyslipidemia can change the intestinal microbiota and affect the reproduction of metabolites and short-chain fatty acids, leading to imbalance of the intestinal microbiota. Therefore, bioactive components decrease the bacteria that cause imbalances and increase beneficial bacteria. *Firmicutes* is more efficient with sugar metabolism than beneficial bacteria, which favors energy absorption and storage, promoting obesity.²⁹

ethanol extract of the pulp for 28 days in mice exposed to cigarette smoke, resulting in the ethanol extract of tree tomato being able to prevent development of atherosclerosis due to oxidative stress.⁴⁵

Salazar-Lugo et al⁴⁶ they studied 54 volunteers with hypercholesterolemia and hypertriglyceridemia and consumed tree tomato juice (100 g of fruit in 150 mL of water) daily for 6 weeks, the result was a significant decrease in total cholesterol and LDL (Figure 6).

The beneficial effects on health include weight reduction, increase in antioxidant enzymes, prevents postprandial hyperglycemia and has anti-inflammatory effects with the reduction of cytokines such as TNF- α and IL-6.⁴³ Likewise, the seed and skin of the fruit showed antioxidant activity by suppressing the generation of free radicals.^{44,49} The crude




Figure 5 Bioactive compounds present in Pineapple (*Ananas comosus*) and their effect in vivo. Created in BioRender. Flores Flores, M. (2025) <https://BioRender.com/r70d826>.


Notes: By inhibiting HMG CoA reductase, statins reduce cholesterol synthesis in the liver. Carnitine palmitoyl transferase I (CPT-I) converts long-chain fatty acyl-CoA molecules into their corresponding acylcarnitine molecules, which is associated with the inhibition of fatty liver formation.³⁷

extract of tree tomato inhibits the activity of the enzyme α -glucosidase,⁴⁷ α -amylase and lipase, and enzymes that are related to metabolic syndrome.⁴⁸

The skin, pulp and seeds of the fruit are rich in fiber, total phenolics, anthocyanins, carotenoids, ascorbic acid,^{44,47,49} condensed and hydrolyzable tannins,⁵⁰ rosmarinic acid and caffeic acid.⁵¹

Coffee (*Coffea*)

Coffee is one of the most consumed beverages worldwide due to its unique flavour and the role it plays in social gatherings; it belongs to the family *Rubiaceae*, subfamily *Cinchonoideae* and tribe *Coffeae* and to the genus *Coffea*.⁵² The pulp and skin of coffee stands out for its content of polyphenols and flavonoids such as chlorogenic acid (CGA),






Figure 7 Bioactive compounds present in Coffee (*Coffea*) and their effect in vivo. Created in BioRender. Flores Flores, M. (2025) <https://BioRender.com/o96y523>.

Notes: The mechanisms of lipid control are probably through the inhibition of intestinal cholesterol absorption and the antioxidant effects of polyphenols such as chlorogenic acid (CGA), epicatechin (EC).⁵³ Additionally, there is evidence linking coffee to a decreased risk of Parkinson's and Alzheimer's diseases.⁵⁷

Among the health benefits, it is found to be the greatest source with antioxidant capacity, cholesterol reduction, anti-inflammatory, diseases that reduce cognitive function,⁵³ and prevention of hepatic steatosis, insulin resistance,⁵⁹ protects against reactive oxygen radicals, reduces the levels of glycosylated hemoglobin (HbA1c),⁵⁷ prevents cancer and degenerative diseases, aging⁶⁰ and reduces the risk of suffering from diabetes mellitus.⁶¹ In other research, it is highlighted that coffee pulp reduces blood glucose, lipid peroxidation, and increases hepatic, renal and cardiac glutathione values. Likewise, it promotes the activity of catalase, glutathione peroxidase and superoxide dismutase in rats induced to diabetes.⁵⁵

Apple (*Malus domestica*)

The apple is the most consumed fruit worldwide, it belongs to the *Rosaceae* family. They are rich in anthocyanins and low in calories.^{51,62} Likewise, the peel is rich in flavonoids (quercetin), polyphenols⁶³ and pectin, a bioactive component of the cell wall that has an effect against cholesterol.⁶⁴

Studies with rats showed that ingesting 114 mg/kg of apple peel extract for 30 days reduces LDL levels and increases HDL.⁶³ The use of 20% of the feed weight with sun-dried apple peels for 43 days significantly reduced triglyceridemia levels⁶⁵ and where 600–900 mg/kg of polyphenolic apple peel extract was used for 8 weeks, increased HDL levels and decreased TC, TG and LDL.⁶⁶ Using 15 mL/kg of apple juice for 21 days decreased levels of TC, TG, LDL and increased HDL.⁶⁷ Forty male Wistar rats received Apple cider vinegar (6 g vinegar/100 g animal food) for 4 weeks, the result was 47% reduction in LDL-c levels and 34% increase in HDL-c levels.

Study in 23 healthy human patients, it was confirmed that consuming 550 g/day of whole apples and 22 g/day of apple pulp for 4–5 weeks reduces TC and LDL levels.⁶⁴ Seventy participants with hyperlipidemia and type 2 diabetes received 20 mL apple vinegar per day using an 8-week, the result showed decreased the dyslipidemia (Figure 8).⁶⁸




Figure 8 Bioactive compounds present in Apple (*Malus domestica*) and their effect in vivo. Created in BioRender. Flores Flores, M. (2025) <https://BioRender.com/t54q725>.
Notes: HMG-CoA reductase (HMG) inhibits cholesterol biosynthesis in the liver before the synthesis of mevalonate.⁶³ It also has an impact on reducing high blood pressure caused by increased cholesterol or triglycerides.⁶²

Its health effects are antioxidant, anti-obesity, anti-hyperlipidemic activity, prevention of metabolic syndrome,^{62,63} lower risk of death from cerebrovascular disease⁶⁹ and changes in the intestinal microbiota of rats that consumed whole apples.⁷⁰ In addition, apple peels have an effect against high blood glucose and insulin levels⁶⁵ and a decrease

in serum biomarkers of inflammation such as IL-6, IL-8 and TNF- α , an effect against cancer and gastrointestinal protection,⁶⁷ the source of acetic acid and phytochemicals (apple vinegar), can improve glycemic indices and reduce starch digestion.⁶⁸

Passion Fruit (*Passiflora edulis*)

Passiflora edulis, commonly referred to as passion fruit or egg fruit, is indigenous to Brazil and a member of the *Passifloraceae* family of passion flowers. Grown in tropical and subtropical climates, it is a perennial herbaceous vine.⁷¹

Passion fruit oils reduce triglyceride, total cholesterol, and low-density lipoprotein cholesterol content in cells (Pancreatic lipase and Cholesterol esterase).⁷¹ Seven obese male *db/db* mice received 7 g of bark of *Passiflora edulis* (BPe) kg of chow corresponding to 1.5 g/kg of body weight for 8 weeks, the result showed reduction of triglycerides and total cholesterol.⁷² Fifteen offspring from treated nondiabetic dams rats and 15 offspring from treated diabetic dams rats consumed at a dose of 0.58 g/kg once a day (early morning) for 30 consecutive days of passion fruit juice (oral gavage), the result showed significantly reduced low-density lipoprotein cholesterol levels, total cholesterol triglyceride and increased high-density lipoprotein cholesterol level.⁷³ Groups of 6 to 13 Male Wistar rats was induced diabetes and treated with pectin daily for 5 days (0.5–25 mg/kg orally), and the result was pectin decreased triglyceride levels in diabetic rats.⁷⁴ Eight rats induced hyperlipidemic consumed 4.2 mL/200gBB/day of purple passion juice for 14 days, the result was a significant decrease in triglyceride levels.⁷⁵

In other study, 43 adults with dyslipidemia consumed 300 mL jelly drink containing polyphenol-rich roselle calyces extract and passion fruit juice with pulp concentrate (RP jelly drink) for 8 weeks of intervention; results showed a significant decrease in LDL-C and TG, respectively.⁷⁶ Twenty participants with dyslipidemia received 250 mL/day of yellow passion fruit peel juice for 15 days, the result show that HDL level was increased and LDL level decreased.⁷⁷ Eighteen patients with HIV lipodystrophy consumed 30 g of passion fruit peel flour daily for 90 days and diet therapy counseling, and the result was effective in reducing total cholesterol, triacylglycerides, LDL-C and HDL-C increased (Figure 9).⁷⁸

Another diseases that passion fruit can help are anxiolytic, antioxidant, antibacterial, antitumor, anti-inflammatory, hypolipidemic properties,⁷¹ epilepsy, headache, abdominal pain, diabetes mellitus,⁷³ fruit peel in cases of asthma,⁷⁵ reduction of malondialdehyde (MDA) and tumor necrosis factor- (TNF-).⁷⁶ Also, can be helping to reduce the obesity and complications in target organs such as liver, heart and kidney.⁷²

Passion fruit seed oils are rich in unsaturated fatty acids (linoleic acid, C18:2), lipid-soluble acids, vitamin E, β -carotene, proteins, minerals, polyphenols, flavonoids, lipids⁷¹ and vitamin C. On the other hand, passionflower bark flour (*Passiflora edulis*) is rich in pectin, a soluble fiber, able to form a gel.⁷² The pulp is composed of polyphenols, alkaloids, flavonoids, tocopherols, carotenoids and ascorbic acid.⁷⁶

Conclusion

This review gives an overview of the benefit of fresh fruits and their various parts as an alternative for reducing dyslipidemia, as well as for the benefit of other types of diseases related to diet, such as the case of some tumors, types of cancer, diarrhea, cardiovascular diseases, diabetes, etc. It should be noted that among the components that are mostly related to the reduction of free radicals, which prevents lipid peroxidation, are phenolic compounds such as anthocyanins, flavonoids and tannins, which fulfill a powerful antioxidant function. On the other hand, this benefit of reducing dyslipidemia is not only provided by soluble fiber (which we usually find in the edible and inedible parts of fruits) but also by polyphenols such as anthocyanins, thiamine and oils found in the seeds of fruits. The various parts of fruits such as leaves, stems, peels, pulp and seeds are used as prebiotics, which can increase beneficial bacteria, which by producing short-chain fatty acids also fulfill the function of preventing dyslipidemia, among other benefits.




Figure 9 Bioactive compounds present in Passion fruit (*Passiflora edulis*) and their effect in vivo. Created in BioRender. Flores flores, M. (2025) <https://BioRender.com/u41t501>.

Notes: Flavonoids can enhance the conversion of cholesterol into bile acids and restrict its absorption, lowering blood cholesterol levels. Beta-carotene can enhance the elimination of cholesterol through feces and decrease its absorption in the intestines. In the colon, this fiber will ferment and transform into short-chain fatty acids. It can make the intestinal contents more viscous, which will cause the cholesterol from the bile salts to be expelled in the feces.⁷¹

Acknowledgments

Graphical abstract and Figures 1-9 are created in BioRender. Flores Flores, M. (2025) <https://BioRender.com/p88y952>. Diyar Wakkas, Faculty of Law, University of Amsterdam, for his collaboration in the English grammar review.

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

This manuscript was funded by the Universidad Nacional de la Amazonía Peruana with a Rectoral Resolution N° 0437-2025-UNAP.

Disclosure

The authors report no conflicts of interest in this work.

References

- Chrigui S, Taieb SH, Jemai H, et al. Anti-obesity and anti-dyslipidemic effects of salicornia arabica decocted extract in tunisian psammomys obesus fed a high-calorie diet. *Foods*. 2023;12(6):1185. doi:10.3390/FOODS12061185
- Berberich AJ, Hegele RA. A modern approach to dyslipidemia. *Endocr Rev*. 2022;43(4):611–653. doi:10.1210/ENDREV/BNAB037
- Fahmy H, Hegazi N, El-Shamy S, Farag MA. Pomegranate juice as a functional food: a comprehensive review of its polyphenols, therapeutic merits, and recent patents. *Food Funct*. 2020;11(7):5768–5781. doi:10.1039/D0FO01251C
- Isas AS, Escobar F, Álvarez-Villamil E, et al. Fermentation of pomegranate juice by lactic acid bacteria and its biological effect on mice fed a high-fat diet. *Food Biosci*. 2023;53:102516. doi:10.1016/j.fbio.2023.102516
- Halim NAA, Karim CF, Mohammed IK, et al. Anti-obesity effect of methanolic extracts of local punica granatum in high-fat diet-induced obese rats. *Malaysian J Med Health Sci*. 2023;18(2):228–236. doi:10.47836/MJMHS.19.2.33
- Rouhi SZT, Sarker MMR, Rahmat A, Alkahtani SA. The effect of pomegranate fresh juice versus pomegranate seed powder on metabolic indices, lipid profile, inflammatory biomarkers, and the histopathology of pancreatic islets of Langerhans in streptozotocin-nicotinamide induced type 2 diabetic Sprague-Dawley rats. *BMC Complement Altern Med*. 2017;17(1):1–13. doi:10.1186/S12906-017-1667-6/TABLES/3
- Haghighian MK, Raftar M, Hemmati S, Haghravan S, Asghari-Jafarabadi M. Effects of pomegranate (*Punica granatum* L.) peel extract supplementation on serum lipid profile and oxidative stress in obese women with knee osteoarthritis: a double blind, randomized, placebo controlled study. *Adv Integr Med*. 2021;8(2):107–113. doi:10.1016/J.AIMED.2020.05.001
- Kojadinovic M, Glibetic M, Vucic V, et al. Short-term consumption of pomegranate juice alleviates some metabolic disturbances in overweight patients with dyslipidemia. *Journal of Medicinal Food*. 2021;24(9):925–933. doi:10.1089/JMF.2020.0122
- Hamoud S, Hayek T, Volkova N, et al. Pomegranate extract (POMx) decreases the atherogenicity of serum and of human monocyte-derived macrophages (HMDM) in simvastatin-treated hypercholesterolemic patients: a double-blinded, placebo-controlled, randomized, prospective pilot study. *Atherosclerosis*. 2014;232(1):204–210. doi:10.1016/j.atherosclerosis.2013.11.037
- Estrada-Luna D, Carreón-Torres E, Bautista-Pérez R, et al. Microencapsulated pomegranate reverts high-density lipoprotein (HDL)-induced endothelial dysfunction and reduces postprandial triglyceridemia in women with acute coronary syndrome. *Nutrients*. 2019;11(8):1710. doi:10.3390/NU11081710
- Abedini M, Ghasemi-Tehrani H, Tarrahi MJ, Amani R. The effect of concentrated pomegranate juice consumption on risk factors of cardiovascular diseases in women with polycystic ovary syndrome: a randomized controlled trial. *Phytother Res*. 2021;35(1):442–451. doi:10.1002/PTR.6820
- Shema-Didi L, Kristal B, Sela S, Geron R, Ore L. Does Pomegranate intake attenuate cardiovascular risk factors in hemodialysis patients? *Nutr J*. 2014;13(1):18. doi:10.1186/1475-2891-13-18
- Grabež M, Škrbić R, Stojiljković MP, et al. A prospective, randomized, double-blind, placebo-controlled trial of polyphenols on the outcomes of inflammatory factors and oxidative stress in patients with type 2 diabetes mellitus. *Rev Cardiovasc Med*. 2022;23(2):57. doi:10.31083/J.RCM2302057/2153-8174-23-2-057/FIG1.JPG
- Mirmiran P, Fazeli MR, Asghari G, Shafiee A, Azizi F. Effect of pomegranate seed oil on hyperlipidaemic subjects: a double-blind placebo-controlled clinical trial. *Br J Nutr*. 2010;104(3):402–406. doi:10.1017/S0007114510000504
- Esmailzadeh A, Tahbaz F, Gaieni I, Alavi-Majd H, Azadbakht L. Concentrated pomegranate juice improves lipid profiles in diabetic patients with hyperlipidemia. *Journal of Medicinal Food*. 2004;7(3):305–308. doi:10.1089/JMF.2004.7.305
- Esmailzadeh A, Tahbaz F, Gaieni I, Alavi-Majd H, Azadbakht L. Cholesterol-lowering effect of concentrated pomegranate juice consumption in type II diabetic patients with hyperlipidemia. *Int J Vitamin Nutrition Res*. 2006;76(3):147–151. doi:10.1024/0300-9831.76.3.147/PDF
- Mahesar SA, Kori AH, Sherazi STH, Kandhro AA, Laghari ZH. Pomegranate (*Punica granatum*) seed oil. *Fruit Oils*. 2019;691–709. doi:10.1007/978-3-030-12473-1_37
- Aviram M, Volkova N, Coleman R, et al. Pomegranate phenolics from the peels, arils, and flowers are antiatherogenic: studies in vivo in atherosclerotic apolipoprotein E-deficient (E0) mice and in vitro in cultured macrophages and lipoproteins. *J Agric Food Chem*. 2008;56(3):1148–1157. doi:10.1021/JF071811Q/SUPPL_FILE/JF071811QSI20071122_080948.PDF
- Aladaileh SH, Saghir SAM, Murugesu K, et al. Antihyperlipidemic and antioxidant effects of Averrhoa carambola extract in high-fat diet-fed rats. *Biomedicine*. 2019;7(3):72. doi:10.3390/BIOMEDICINES7030072
- Saghir S AM, Sadikun A, Al-Suede F SR, Majid A MSA, Murugaiyah V. Antihyperlipidemic, antioxidant and cytotoxic activities of methanolic and aqueous extracts of different parts of star fruit. *Curr Pharm Biotechnol*. 2016;17(10):915–925. doi:10.2174/1389201017666160603013434
- Abduh MS, Saghir SAM, Al Hroob AM, et al. Averrhoa carambola leaves prevent dyslipidemia and oxidative stress in a rat model of poloxamer-407-induced acute hyperlipidemia. *Front Pharmacol*. 2023;14:198. doi:10.3389/FPHAR.2023.1134812/BIBTEX
- Ramadan NS, El-Sayed NH, El-Toumy SA, et al. Anti-obesity evaluation of Averrhoa carambola L. leaves and assessment of its polyphenols as potential α-glucosidase inhibitors. *Molecules*. 2022;27(16):5159. doi:10.3390/molecules27165159
- Qin L, Zhang X, Zhou X, et al. Protective effect of benzoquinone isolated from the roots of Averrhoa carambola L. on streptozotocin-induced diabetic mice by inhibiting the TLR4/NF-κB signaling pathway. *Diabetes Metab Syndr Obes*. 2020;13:2129. doi:10.2147/DMSO.S241998
- Herman-Lara E, Elvira-Torales LI, Rodríguez-Miranda J, et al. Impact of micronized starfruit (Averrhoa carambola L.) fiber concentrate on lipid metabolism in mice. *Int J Food Sci Nutr*. 2014;65(7):862–867. doi:10.3109/09637486.2014.918590

25. Song H, Zhang Y, Huang Q, Deng R, Zheng X. Averrhoa carambola L. fruit polyphenols ameliorate hyperlipidemia, hepatic steatosis, and hyperglycemia by modulating lipid and glucose metabolism in mice with obesity. *J Sci Food Agric*. 2023;103(13):6531–6539. doi:10.1002/JSSA.12731
26. Shui G, Leong LP. Analysis of polyphenolic antioxidants in star fruit using liquid chromatography and mass spectrometry. *J Chromatogr A*. 2004;1022(1–2):67–75. doi:10.1016/J.CHROMA.2003.09.055
27. Narayana CK. Phytochemicals in fruits and their therapeutic properties. (Taylor & Francis, ed.). 2021. Available from: https://www.google.com.pe/books/edition/Phytochemicals_in_Fruits_and_their_Thera/woFGEAAQBAJ?hl=es&gbpv=0. Accessed April 24, 2025.
28. Li J, Zhang J, Zhang Y, et al. Effect and correlation of Rosa roxburghii trutt fruit vinegar on obesity, dyslipidemia and intestinal microbiota disorder in high-fat diet mice. *Foods*. 2022;11(24):4108. doi:10.3390/FOODS11244108
29. Ni HY, Yu L, Zhao XL, et al. Seed oil of Rosa roxburghii Tratt against non-alcoholic fatty liver disease in vivo and in vitro through PPAR α /PGC-1 α -mediated mitochondrial oxidative metabolism. *Phytomedicine*. 2022;98:153919. doi:10.1016/J.PHYMED.2021.153919
30. Wang L, Li C, Huang Q, Fu X. Polysaccharide from Rosa roxburghii trutt fruit attenuates hyperglycemia and hyperlipidemia and regulates colon microbiota in diabetic db/db mice. *J Agric Food Chem*. 2020;68(1):147–159. doi:10.1021/ACS.JAFC.9B06247/SUPPL_FILE/JF9B06247_SI_001.PDF
31. Ji J, Zhang S, Yuan M, et al. Fermented Rosa roxburghii trutt juice alleviates high-fat diet-induced hyperlipidemia in rats by modulating gut microbiota and metabolites. *Front Pharmacol*. 2022;13. doi:10.3389/FPHAR.2022.883629/FULL
32. Wu PH, Han SCH, Wu MH. Beneficial effects of hydroalcoholic extract from Rosa roxburghii trutt fruit on hyperlipidemia in high-fat-fed rats. *Acta Cardiol Sin*. 2020;36(2):148–159. doi:10.6515/ACS.202003_36(2).20190709A
33. Jin H, Zhang JY, Cai ZY, et al. Therapeutic effects of chigitazar combined with Rosa roxburghii Tratt. in inpatients with antipsychotic-induced metabolic syndrome. *Pak J Pharm Sci*. 2024;37(6):1343. doi:10.36721/PJPS.2024.37.6.REG.1343-1349.1
34. Chen G, Kan J. Characterization of a novel polysaccharide isolated from Rosa roxburghii trutt fruit and assessment of its antioxidant in vitro and in vivo. *Int J Biol Macromol*. 2018;107(PartA):166–174. doi:10.1016/J.IJBIOMAC.2017.08.160
35. Wang H, Li Y, Ren Z, et al. Optimization of the microwave-assisted enzymatic extraction of Rosa roxburghii trutt. polysaccharides using response surface methodology and its antioxidant and α -d-glucosidase inhibitory activity. *Int J Biol Macromol*. 2018;112:473–482. doi:10.1016/J.IJBIOMAC.2018.02.003
36. El-Demerdash FM, Hussien DM, Ghanem NF, AL-Farga AM. Bromelain modulates liver injury, hematological, molecular, and biochemical perturbations induced by aluminum via oxidative stress inhibition. *Biomed Res Int*. 2022;2022(1):1–12. doi:10.1155/2022/5342559
37. Xie W, Zhang S, Lei F, Ouyang X, Du L. Ananas comosus L. Leaf phenols and p-coumaric acid regulate liver fat metabolism by upregulating CPT-1 expression. *Evid Based Complement Alternat Med*. 2014;2014(1). doi:10.1155/2014/903258
38. Seenak P, Kumphune S, Malakul W, Chotima R, Nernpermpisooth N. Pineapple consumption reduced cardiac oxidative stress and inflammation in high cholesterol diet-fed rats. *Nutr Metab*. 2021;18(1):1–10. doi:10.1186/S12986-021-00566-Z/FIGURES/5
39. Namwong A, Kumphune S, seenak P, Chotima R, Nernpermpisooth N, Malakul W. Pineapple fruit improves vascular endothelial dysfunction, hepatic steatosis, and cholesterol metabolism in rats fed a high-cholesterol diet. *Food Funct*. 2022;13(19):9988–9998. doi:10.1039/D2FO01199A
40. El-Shazly SA, Ahmed MM, AL-Harbi MS, Alkafafy ME, El-Sawy HB, Amer SAM. Physiological and molecular study on the anti-obesity effects of pineapple (Ananas comosus) juice in male Wistar rat. *Food Sci Biotechnol*. 2018;27(5):1429. doi:10.1007/S10068-018-0378-1
41. Xie W, Xing D, Sun H, Wang W, Ding Y, Du L. The effects of ananas comosus L. Leaves on diabetic-dyslipidemic rats induced by alloxan and a high-fat/high-cholesterol diet. *Am J Chin Med*. 2012;33(1):95–105. doi:10.1142/S0192415X05002692
42. Chen CH, Hsia CC, Hu PA, et al. Bromelain ameliorates atherosclerosis by activating the TFEB-mediated autophagy and antioxidant pathways. *Antioxidants*. 2023;12(1). doi:10.3390/ANTIOX12010072
43. Abdul Kadir NAA, Rahmat A, Jaafar HZE. Protective effects of tamarillo (Cyphomandra betacea) extract against high fat diet induced obesity in Sprague-Dawley rats. *J Obes*. 2015;2015:1–8. doi:10.1155/2015/846041
44. Orqueda ME, Rivas M, Zampini IC, et al. Chemical and functional characterization of seed, pulp and skin powder from chilito (Solanum betaceum), an Argentine native fruit. Phenolic fractions affect key enzymes involved in metabolic syndrome and oxidative stress. *Food Chem*. 2017;216:70–79. doi:10.1016/j.foodchem.2016.08.015
45. Khaerunnisa S, Hidayati HB, Susanto J, Setiawati Y, Suhartati S. Atheroprotective effect of Solanum betaceum on rat exposed to cigarette smoke. *Int J Appl Pharm*. 2019;11(Special Issue 5):145–149. doi:10.22159/IJAP.2019.V11S5.T1005
46. Salazar-Lugo R, Barahona A, Ortiz K, et al. Effect of consumption of tree tomato juice (Cyphomandra betacea) on lipid profile and glucose concentrations in adults with hyperlipidemia, Ecuador. *Arch Latinoam Nutr*. 2016;66(2):121–128.
47. Puspawati GAKD, Marsono Y, Armunanto R, Supriyadi S. Inhibitory potency of Indonesian tamarillo (Solanum betaceum Cav) crude extract against α -glucosidase enzyme activity. *Curr Res Nutr Food Sci*. 2018;6(2):392–403. doi:10.12944/CRNFSJ.6.2.14
48. Orqueda ME, Torres S, Zampini IC, et al. Integral use of Argentinean Solanum betaceum red fruits as functional food ingredient to prevent metabolic syndrome: effect of in vitro simulated gastroduodenal digestion. *Heliyon*. 2020;6(2):e03387. doi:10.1016/j.heliyon.2020.e03387
49. Orqueda ME, Torres S, Verón H, et al. Physicochemical, microbiological, functional and sensory properties of frozen pulp of orange and orange-red chilito (Solanum betaceum Cav.) fruits. *Sci Hortic*. 2021;276:109736. doi:10.1016/J.SCIENTA.2020.109736
50. Correa Uriburu FM, Zampini IC, Maldonado LM, Gómez Mattson M, Salvatori D, Isla MI. Powdered beverage from native plants from Argentina (Zuccagnia punctata and Solanum betaceum) obtained by spray-drying: a promising source of antioxidant compounds. *Plants*. 2023;12(8):1646. doi:10.3390/plants12081646
51. Moreno MA, Orqueda ME, Gómez-Mascaraque LG, Isla MI, López-Rubio A. Crosslinked electrospun zein-based food packaging coatings containing bioactive chilito fruit extracts. *Food Hydrocoll*. 2019;95:496–505. doi:10.1016/J.FOODHYD.2019.05.001
52. Murthy PS, Madhava Naidu M. Sustainable management of coffee industry by-products and value addition—A review. *Resour Conserv Recycl*. 2012;66:45–58. doi:10.1016/J.RESCONREC.2012.06.005
53. Rungraung N, Muangpracha N, Trachootham D. Twelve-week safety and potential lipid control efficacy of coffee cherry pulp juice concentrate in healthy volunteers. *Nutrients*. 2023;15(7):1602. doi:10.3390/NU15071602
54. Viñas Meneses M, Gruschwitz M, Schweiggert RM, et al. Identification of phenolic and carotenoid compounds in coffee (Coffea arabica) pulp, peels and mucilage by HPLC electrospray ionization mass spectrometry. In: *24th International Conference on Coffee Science (pp127-135) Costa Rica: Book of Proceedings*; 2018. <https://kerwa.ucr.ac.cr/handle/10669/75350>. Accessed April 24, 2025.

55. Ontawong A, Duangjai A, Muanprasat C, et al. Lipid-lowering effects of *Coffea arabica* pulp aqueous extract in Caco-2 cells and hypercholesterolemic rats. *Phytomedicine*. 2019;52:187–197. doi:10.1016/J.PHYMED.2018.06.021
56. Chang CW, Hsu YJ, Chen YM, Huang WC, Huang CC, Hsu MC. Effects of combined extract of cocoa, coffee, green tea and garcinia on lipid profiles, glycaemic markers and inflammatory responses in hamsters. *BMC Complement Altern Med*. 2015;15(1):269. doi:10.1186/S12906-015-0806-1
57. Shaposhnikov S, Hatzold T, El yamani N, et al. Coffee and oxidative stress: a human intervention study. *Eur J Nutr*. 2018;57(2):533–544. doi:10.1007/S00394-016-1336-4/TABLES/10
58. Sarriá B, Martínez-López S, Sierra-Cinos JL, García-Diz L, Mateos R, Bravo-Clemente L. Regularly consuming a green/roasted coffee blend reduces the risk of metabolic syndrome. *Eur J Nutr*. 2018;57(1):269–278. doi:10.1007/S00394-016-1316-8/METRICS
59. Ontawong A, Boonphang O, Pasachan T, et al. Hepatoprotective effect of coffee pulp aqueous extract combined with simvastatin against hepatic steatosis in high-fat diet-induced obese rats. *J Funct Foods*. 2019;54:568–577. doi:10.1016/J.JFF.2019.02.011
60. Bichler J, Cavin C, Simic T, et al. Coffee consumption protects human lymphocytes against oxidative and 3-amino-1-methyl-5H-pyrido[4,3-b]indole acetate (Trp-P-2) induced DNA-damage: results of an experimental study with human volunteers. *Food and Chemical Toxicology*. 2007;45(8):1428–1436. doi:10.1016/J.FCT.2007.02.001
61. Du Y, Lv Y, Zha W, Hong X, Luo Q. Effect of coffee consumption on dyslipidemia: a meta-analysis of randomized controlled trials. *Nutr Metab Cardiovasc Dis*. 2020;30(12):2159–2170. doi:10.1016/J.NUMECD.2020.08.017
62. Popiolek-Kalisz J, Glibowski P. Apple peel supplementation potential in metabolic syndrome prevention. *Life*. 2023;13(3):753. doi:10.3390/LIFE13030753
63. Susilowati R, Jannah J, Maghfuroh Z, Kusuma M. Antihyperlipidemic effects of apple peel extract in high-fat diet-induced hyperlipidemic rats. *J Adv Pharm Technol Res*. 2020;11(3):128. doi:10.4103/JAPTR.JAPTR_28_20
64. Ravn-Haren G, Dragsted LO, Buch-Andersen T, et al. Intake of whole apples or clear apple juice has contrasting effects on plasma lipids in healthy volunteers. *Eur J Nutr*. 2013;52(8):1875–1889. doi:10.1007/S00394-012-0489-Z/METRICS
65. Gonzalez J, Donoso W, Sandoval N, et al. Apple peel supplemented diet reduces parameters of metabolic syndrome and atherogenic progression in ApoE^{-/-} mice. *Evid Based Complement Alternat Med*. 2015;2015:1–10. doi:10.1155/2015/918384
66. Jia M, Ren D, Nie Y, Yang X. Beneficial effects of apple peel polyphenols on vascular endothelial dysfunction and liver injury in high choline-fed mice. *Food Funct*. 2017;8(3):1282–1292. doi:10.1039/C7FO00147A
67. Fathy SM, Drees EA, Mondégo-Oliveira R. Protective effects of Egyptian cloudy apple juice and apple peel extract on lipid peroxidation, antioxidant enzymes and inflammatory status in diabetic rat pancreas. *BMC Complement Altern Med*. 2016;16(1):1–14. doi:10.1186/S12906-015-0957-0/FIGURES/5
68. Gheflati A, Bashiri R, Ghadiri-Anari A, Reza JZ, Kord MT, Nadjarzadeh A. The effect of apple vinegar consumption on glycemic indices, blood pressure, oxidative stress, and homocysteine in patients with type 2 diabetes and dyslipidemia: a randomized controlled clinical trial. *Clin Nutr ESPEN*. 2019;33:132–138. doi:10.1016/J.CLNESP.2019.06.006/ATTACHMENT/550A3114-8897-4B2C-8FDD-28A2526CCA0D/MMC1.XML
69. Hodgson JM, Prince RL, Woodman RJ, et al. Apple intake is inversely associated with all-cause and disease-specific mortality in elderly women. *Br J Nutr*. 2016;115(5):860–867. doi:10.1017/S0007114515005231
70. Licht TR, Hansen M, Bergström A, et al. Effects of apples and specific apple components on the cecal environment of conventional rats: role of apple pectin. *BMC Microbiol*. 2010;10(1):1–11. doi:10.1186/1471-2180-10-13/TABLES/6
71. Chen P, Zhang X, Yang Q, et al. Extraction and characterization of passion fruit seed oil and investigation of its hypolipidemic activity. *J Agric Food Res*. 2025;19:101672. doi:10.1016/J.JAFR.2025.101672
72. Panelli MF, Pierine DT, de Souza SLB, et al. Bark of *Passiflora edulis* treatment stimulates antioxidant capacity, and reduces dyslipidemia and body fat in db/db mice. *Antioxidants*. 2018;7(9):120. doi:10.3390/ANTIOX7090120
73. Barbalho SM, Damasceno DC, Spada APMH, et al. Effects of *Passiflora edulis* on the metabolic profile of diabetic Wistar rat offspring. *Journal of Medicinal Food*. 2011;14(12):1490–1495. doi:10.1089/JMF.2010.0318
74. Silva DC, Freitas ALP, Pessoa CDS, et al. Pectin from *Passiflora edulis* shows anti-inflammatory action as well as hypoglycemic and hypotriglyceridemic properties in diabetic rats. *Journal of Medicinal Food*. 2011;14(10):1118–1126. doi:10.1089/JMF.2010.0220
75. Muntafiah A, Pratama TS, Rachmawati A, Wulandari DW, Santosa Q. Comparison of purple passion juice (*Passiflora edulis* var. *edulis*) and simvastatin on lipid-lowering effect of hyperlipidemic rats model. In: *Proceedings of the 1st Jenderal Soedirman International Medical Conference in conjunction with the 5th Annual Scientific Meeting (Temilnas) Consortium of Biomedical Science Indonesia*; 2020:148–154. doi:10.5220/0010489301480154.
76. Khongrum J, Yingthongchai P, Boonyapranai K, et al. Antidyslipidemic, antioxidant, and anti-inflammatory effects of jelly drink containing polyphenol-rich rosele calyces extract and passion fruit juice with pulp in adults with dyslipidemia: a randomized, double-blind, placebo-controlled trial. *Oxid Med Cell Longev*. 2022;2022:4631983. doi:10.1155/2022/4631983
77. Adi DI. The effect of yellow passion fruit peel juice (*Passiflora edulis* f. *flavicarpa* deg.) on HDL and LDL levels in type 2 diabetes mellitus patients in the working area of teppo health centre pinrang regency. *Indian J Public Health Res Dev*. 2019;10(4):984–989. doi:10.5958/0976-5506.2019.00836.2
78. Marques S Do SF, Libonati RMF, Sabaa-Srur AUO, et al. Evaluation of the effects of passion fruit peel flour (*Passiflora edulis* fo. *flavicarpa*) on metabolic changes in HIV patients with lipodystrophy syndrome secondary to antiretroviral therapy. *Rev Bras Farmacogn*. 2016;26(4):420–426. doi:10.1016/J.BJP.2016.03.002

Vascular Health and Risk Management**Publish your work in this journal**

Vascular Health and Risk Management is an international, peer-reviewed journal of therapeutics and risk management, focusing on concise rapid reporting of clinical studies on the processes involved in the maintenance of vascular health; the monitoring, prevention and treatment of vascular disease and its sequelae; and the involvement of metabolic disorders, particularly diabetes. This journal is indexed on PubMed Central and MedLine. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/vascular-health-and-risk-management-journal>

Dovepress
Taylor & Francis Group