

Assessment of Government Intervention Effectiveness in Scrubbing COVID-19 Pandemic and Its Pros and Cons on Educational Activities, the Case of Dire Dawa City

Gizaw Tesfaye Yifru¹
Bekele Yazie Derso²
Alemayehu Zewdie Shibeshi³

¹College of Natural and Computational Sciences, Dire Dawa University, Dire Dawa, Ethiopia; ²College of Medicine and Health Sciences, Dire Dawa University, Dire Dawa, Ethiopia; ³College of Social Science and Humanities, Dire Dawa University, Dire Dawa, Ethiopia

Background: Coronaviruses are known as a large family of viruses known to cause respiratory infections which are considered as a pandemic by WHO and widely distributed all over the globe; causing several damages to all aspects of human being's life.

Objective: To assess and identify the determinants of government intervention effectiveness in scrubbing COVID-19 and its pros and cons on educational activity in Dire Dawa City.

Methods: Stratified random sampling was hired to draw a sample of teachers from a list of teachers that were taught at both selected private and public educational institutes.

Results: A 57.6% of educators responded as the government was effective in mitigating the pandemic. Out of 250 educators, 9.6%, 10.8%, 8.4%, 38.4%, and 32.8% were reported that they have very poor, poor, neutral, good, and very good level of knowledge about COVID-19, respectively. The respondents' perceptions about the pandemic were rated as not very fear inducing, not fear inducing, neutral, fear inducing, and very fear inducing 3.2%, 6.4%, 6.8%, 55.2% and 28.4%, respectively. Logistic regression indicates nine predictors were significant: avoiding touching one's body with unwashed hands, disinfecting surfaces, fatigue, getting flu vaccine, individual limitation in cooperating to cease the disease, how to maintain one's mental health during the isolation, and washing for at least 20 seconds, family care, and self-isolation.

Pros: Reading, family care, watching movies and physical exercise were importance brought by the pandemic to the educators.

Cons: Teachers are obligated to stay at home, unemployment, stress, unable to conduct professional activities, cancellation of training, and loss of motivation.

Conclusion: The results implied us eighty-four variables were significantly associated with government intervention effectiveness, and nine predictors were significantly related with the government's intervention effectiveness in halting the pandemic from logistic regression model.

Keywords: government, teachers, interventions, effectiveness, COVID-19 pandemic, schools, physical distancing, stay at home, educational activities, pros and cons, quarantine

Introduction

Coronaviruses are a large family of viruses known to cause respiratory infections. These can range from the common cold to more serious diseases such as Severe Acute Respiratory Syndrome (SARS) and Middle East Respiratory Syndrome (MERS).¹

Correspondence: Gizaw Tesfaye Yifru
Tel +251912233221
Email gtgizsky@gmail.com

Received: 10 June 2021
Accepted: 30 August 2021
Published: 21 September 2021

Journal of Multidisciplinary Healthcare 2021:14 2669–2693

2669



© 2021 Tesfaye Yifru et al. This work is published and licensed by Dove Medical Press Limited. The full terms of this license are available at <https://www.dovepress.com/terms.php> and incorporate the Creative Commons Attribution – Non Commercial (unported, v3.0) License (<http://creativecommons.org/licenses/by-nc/3.0/>). By accessing the work you hereby accept the Terms. Non-commercial uses of the work are permitted without any further permission from Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial use of this work, please see paragraphs 4.2 and 5 of our Terms (<https://www.dovepress.com/terms.php>).

This new coronavirus originated in Hubei Province, China and the disease caused by the virus is named COVID-19.¹

COVID-19 is most likely to spread from person-to-person through: close contact with a person while they are infectious or in the 24 hours before their symptoms appeared, close contact with a person with a confirmed infection who coughs or sneezes, and touching objects or surfaces contaminated from a cough or sneeze from a person with a confirmed infection, and then touching your mouth or face.¹

Epidemiological Situation of Novel Corona Virus Pandemic

Globally, as of 4:38pm CEST, 25 August 2021, there have been 213,050,725 confirmed cases of COVID-19, including 4,448,352 deaths, reported to WHO² and as of 22 August 2021, a total of 4,619,976,274 vaccine doses have been administered.²

In Africa: 5,539,401 confirmed cases and in Ethiopia, from 3 January 2020 to 4:38pm CEST, 25 August 2021, there have been 297,997 confirmed cases of COVID-19 with 4580 deaths, reported to WHO.²

As of 22 August 2021, a total of 2,343,609 vaccine doses have been administered.² Whereas, in Dire Dawa City, as of August 25, there have been 5374 confirmed cases of COVID-19 and 86 deaths reported.³

Coronavirus pandemic is spreading globally as of April, 2020: 2,994,690 total confirmed cases, 207,270 deaths whereas in the USA: 965,933 total number of confirmed cases, and 54,877 total deaths reported;⁴ whereas in Ethiopia 123 cases, 3 deaths were reported;⁵ and in Dire Dawa City 9 cases and 0 deaths were reported.³

In Addis Ababa and Dire-Dawa City Administrations, multi-sectoral steering committees were established to coordinate COVID-19.⁶ Measures were taken including banning mass gathering and crowded public in transport. Humanitarian impact, established multi-sectoral coordination meetings disrupted.

Local rehabilitation centers have been impacted. Actions taken: volunteers mobilized to create awareness and resource mobilization; market regulation and stabilization measures; provide sanitation and hygiene materials.

The Negative Effects of COVID-19 on Health Professionals

Zhang et al⁷ reported that their findings provide evidence to enable healthcare organizations to identify staff concerned about job satisfaction, life satisfaction, and turnover intention to enable early actions so that these staffs can remain motivated to fight the prolonged COVID-19 pandemic. Another study done by Zhang et al⁸ implied that healthcare workers are facing high workloads with resource constraints and risk of virus exposure, and healthcare organizations need to support their healthcare workers to reduce their anxiety. Additionally, the study conducted by Yáñez et al⁹ showed that Peru's healthcare workers' anxiety and mental distress decreased as the distance from the epicenter increases, corroborating the ripple effect and disconfirming the typhoon eye theory.

Yáñez et al⁹ have reported that their results can help guide mental health service providers toward vulnerable groups of healthcare workers that are closer to Lima, the COVID-19 epicenter in Peru.

Further, Alvarez-Risco et al¹⁰ reported that Peru was forced into telemedicine due to the pandemic, and the government is working to improve Internet coverage.

Negative Effects of the Pandemic on Citizens

According to the study finding done by Alvarez-Risco et al¹¹ showed that multitasking behavior was found to negatively influence self-efficacy of -0.332 , whereas self-efficacy showed a positive influence of 0.325 on academic performance.

Alvarez-Risco et al¹² reported that the pandemic has changed the world, creating the need for new actions from society, including universities and companies.

A study done by Aleidi et al¹³ implied that self-medication impacts both negatively and positively the health of people, which has become evident during the COVID-19 pandemic. Aleidi et al¹³ suggested that a continuous awareness and sensitization about the risks of self-medication are warranted.

Further, a study performed by Alvarez-Risco et al¹⁴ signalled that the COVID-19 outbreak has highlighted the need to target infodemics that can be as detrimental as an actual epidemic.

Furthermore, Yáñez et al¹⁵ reported that we must be cautious in Peru, where there is a precarious health system, cases keep increasing in small children, pediatric hospitals are saturated with adult patients, and public policies are not necessarily following the global epidemiological alerts.

Negative Effects of the Outbreak on Firms

A research done by Yan et al¹⁶ implied that hospitality industry worldwide is suffering under the COVID-19 pandemic.

On the other hand, the study conducted by Alvarez-Risco et al¹⁷ showed that the impacts of education development, conceptual development, and country supports on the entrepreneur's ability to carry out green entrepreneurship were positive.

Eventually, the Ethiopian government intervenes to take several actions in order to scrub the outbreak of COVID-19. But, the question is, are these all government interventions effective or not?

Thus, the effectiveness of these interventions can be measured via indicator variables. Alternatively, if the interventions taken by the government were effective, they have to meet the pre-specified goals: reduce spread of the disease, reduce death, increase recovery rate, increase the probability of finding the person who has been close contact with patient via tracing task force, increased awareness of both rural and urban society, acceptance and practical implementations of stay at home policy, physical distancing, no handshake, frequent hand wash with soap and water, helping the poor, and others.

The happening of multi-dimensionality impact of COVID-19 outbreak worldwide is obvious that our investigation was intended to identify government interventions effectiveness to scrubbing COVID-19 in Dire Dawa City and to find out determinants of government intervention effectiveness, as well as the specific impact of the disease on educational sector in the city, was studied.

Significance of the Study

The primary importance of the study was to provide information for government and policymakers. Secondly, it is worthy for Dire Dawa City as means of finding out the advantage and the disadvantage of the pandemic on its educational activities and it can be used as remedial action to alleviate from these challenges and it may also use to

reduce the possible damage that might face the city in the future due to this wide spreading disaster. Finally, it can be used to identify the determinants of government intervention effectiveness in fighting the disease in the city. Further, it can be used as an input for the study that would be conducted in the area.

Materials and Methods

Study Area and Period

Dire Dawa City was our study area, found 515 K.M. far away to the East of Ethiopia's capital city. The survey was conducted in the time interval of May 01, and June 01, 2020.

Study Design and Population

A cross-sectional study design held used to generate quantitative data. The parent population of the study is all teachers (936) who are being teaching in Dire Dawa City selected private and public educational institutions in the year 2020.

Inclusion criteria: all teachers who teach in the selected schools and that are permanently residing in the city and those were on duty considered.

Exclusion criteria: teachers who teach in rural and who did no found on duty omitted.

Sample Size and Sampling Technique

To calculate the sample size, level of significance $\alpha = 0.05$, allowable margin of error ($d = 0.04$), and $P = 50\%$, $Q = 50\%$ was employed. The sample was determined to be 366. Stratified random sampling technique was used. Thus, the study population was categorized into three broad strata on the bases of school type as stratification factor: primary schools, secondary schools and tertiary-level institutes. In order to determine the sample size of each stratum, proportional allocation technique was employed, then the simple random sample of study participants was drawn from each stratum.

Data Collection, Instruments and Procedures

To generate the data from the primary sources, interviewers administered structured questionnaire. The questionnaire was adopted from World Health Organization (WHO). Semi-telephone survey was used to collect household data pertaining to knowledge, perception, attitude, and opinion about the pandemic, symptoms of the

outbreak, prevention methods of COVID-19, variables related to government effectiveness in responding and managing the pandemic, education professionals related and educational institute-related variables.

Data Quality Control

In order to minimize the errors in data collection, the enumerators were trained regarding the objectives of the investigation and the content of the research tool. Supervisors have monitored the enumerators to validate the collected data. The data quality was guaranteed via statistical software.

Measurement and Definitions

Generally, we have measured the effectiveness of the government intervention action towards fighting COVID-19 as (effective =1, not effective =0). Independent variables include interventions taken by the government of Ethiopia and recommended by the World Health Organization – no handshake, social or physical distancing (applicable, not applicable), and lockdown (appropriate, not appropriate).

Data Processing and Analysis

SPSS26, R, and R Studio softwares have applied for data validation and data analysis. Proportions and other measures which describe the data and Inferential Statistics employed for data analysis. The results are presented via [Tables 1–17](#) and [figure3](#).

Ethical Consideration

Dire Dawa University research ethics review committee approved there were no ethical issues. Additionally, support letter with this was given to Dire Dawa Administration Education and Health Bureaus, and permission from each School's Principals and Administrative organ of the Educational Institutes involved in the study has found. After a detailed explanation of the study objectives, all participants were voluntary to be interviewed about the topic under the consideration; informed oral consent was sought before the interview. This was done due to the pandemic season and the data collection method employed. Secret was kept by excluding names, and confidentiality was guaranteed with a considerable amount of care. The information gained was kept anonymous to all of the interviewees. The study was done as per the Declaration of Helsinki requirements.

Results

Descriptive Statistics Analysis Result

The primary aim of the study was dealt with government's effectiveness in scrubbing the spread of COVID-19 pandemic in Dire Dawa City. Accordingly, out of 366 total teachers 250 of them were participated in the study and 57.6% of them responded as the government was effective in ceasing the pandemic.

To start with School types of study respondents, 38.0% of them were reported as they are teaching in Primary Schools, 31.6% of them teach in Secondary Schools, and 30.4% of them teach in Technical colleges and Universities. Out of 250 educators, 9.6%, 10.8%, 8.4%, 38.4%, and 32.8% were reported that they have very poor, poor, neutral, good, and very good level of knowledge about COVID-19, respectively.

The respondents' perception about the pandemic were rated as: not very fear inducing, not fear inducing, neutral, fear inducing, and very fear inducing 3.2%, 6.4%, 6.8%, 55.2% and 28.4% about pandemic, respectively.

Bivariate Analysis Result

This section reports the association between the effectiveness of government intervention measures taken to cease the pandemic each of the predictor variables that was significant.

The proportions of educators who rate their level of knowledge about the pandemic to fall in the categories of very poor, poor, neutral, good, and very good were 83.3%, 55.6%, 52.4%, 43.8%, and 68.3%, respectively, were responded as the government intervention measure was effective in halting the pandemic.

This implies that the proportion of survey respondents' rate of government intervention effectiveness decreases as their level of knowledge about the pandemic increases from poor to good and increases for every good level of respondents' knowledge about the outbreak (see [Table 1](#)).

Similarly, the proportion of government effectiveness increases with respondents' knowledge of the symptom of the outbreak. Only 30.0% of study participants were responded as Fever is not among the symptoms of COVID-19 pandemic and rate the government intervention measures taken to scrub the pandemic was effective as compared to 55.8% of who did say that it's among the symptoms of the pandemic and rated that the intervention was effective (see [Table 2](#)).

Table 1 Test of Association Between the Effectiveness of Government Intervention Measures Taken to Cease COVID-19 Pandemic in Dire Dawa City in the Year 2020 and Indicator Variables

Indicator Variables	Respondents' Response Categories	Government Intervention Effectiveness to Cease COVID-19 in DDC						Chi-Square (Sig.)
		Not Effective		Effective		Total		
		Count	%	Count	%	Count	%	
Respondents level of knowledge about the pandemic	Very Poor	4	16.7%	20	83.3%	24	9.6%	18.167 (0.001)
	Poor	12	44.4%	15	55.6%	27	10.8%	
	Neutral	10	47.6%	11	52.4%	21	8.4%	
	Good	54	56.2%	42	43.8%	96	38.4%	
	Very good	26	31.7%	56	68.3%	82	32.8%	
Respondents Susceptibility to be infected with the novel coronavirus	Very unsusceptible	6	30.0%	14	70.0%	20	8.0%	10.363 (0.035)
	Unsusceptible	22	30.6%	50	69.4%	72	28.8%	
	Neutral	30	50.0%	30	50.0%	60	24.0%	
	Susceptible	47	50.5%	46	49.5%	93	37.2%	
	Very Susceptible	1	20.0%	4	80.0%	5	2.0%	
Illness severity due to the disease if happen	Not Very Sever	6	33.3%	12	66.7%	18	7.2%	18.102 (0.001)
	Not Sever	20	57.1%	15	42.9%	35	14.0%	
	Neutral	40	31.0%	89	69.0%	129	51.6%	
	Sever	28	58.3%	20	41.7%	48	19.2%	
	Very Sever	12	60.0%	8	40.0%	20	8.0%	
Knowledge how to protect from coronavirus	I do not know so much	3	16.7%	15	83.3%	18	7.2%	19.511 (0.001)
	I do not know	0	0.0%	2	100.0%	2	0.8%	
	Neutral	4	14.3%	24	85.7%	28	11.2%	
	I know	80	47.9%	87	52.1%	167	66.8%	
	I know so much	19	54.3%	16	45.7%	35	14.0%	
Avoiding an infection with the novel coronavirus in the current situation	Very difficult	29	28.4%	73	71.6%	102	40.8%	16.043 (0.003)
	Difficult	47	54.0%	40	46.0%	87	34.8%	
	Neutral	9	42.9%	12	57.1%	21	8.4%	
	Easy	13	46.4%	15	53.6%	28	11.2%	
	Very easy	8	66.7%	4	33.3%	12	4.8%	

(Continued)

Table 1 (Continued).

Indicator Variables	Respondents' Response Categories	Government Intervention Effectiveness to Cease COVID-19 in DDC						Chi-Square (Sig.)
		Not Effective		Effective		Total		
		Count	%	Count	%	Count	%	
Knowledge and Self-assessed adherence to preventive measures	Not follow at all Not follow Neutral Follow Follow much	3	42.9%	4	57.1%	7	2.8%	18.455 (0.001)
		1	33.3%	2	66.7%	3	1.2%	
		10	47.6%	11	52.4%	21	8.4%	
		60	34.7%	113	65.3%	173	69.2%	
		32	69.6%	14	30.4%	46	18.4%	
Feeling of respondents about the Closeness of COVID-19	Very close to me Close to me Neutral Far away from me Very far away from me	16	51.6%	15	48.4%	31	12.4%	11.584 (0.021)
		48	35.8%	86	64.2%	134	53.6%	
		13	35.1%	24	64.9%	37	14.8%	
		27	62.8%	16	37.2%	43	17.2%	
		2	40.0%	3	60.0%	5	2%	
Newness of the outbreak	Very new to me New to me Neutral Old to me Very old to me	16	38.1%	26	61.9%	42	16.8%	9.907 (0.042)
		59	48.4%	63	51.6%	122	48.8%	
		12	48.0%	13	52.0%	25	10.0%	
		10	50.0%	10	50.0%	20	8.0%	
		9	22.0%	32	78.0%	41	16.4%	
Spread ability of the pandemic	Spread very slowly Spread Slowly Neutral Spread fast Spread very fast	3	33.3%	6	66.7%	9	3.6%	14.754 (0.005)
		51	56.7%	39	43.3%	90	36.0%	
		5	62.5%	3	37.5%	8	3.2%	
		26	31.0%	58	69.0%	84	33.6%	
		21	35.6%	38	64.4%	59	23.6%	
Stressfulness of the disease	Very Stressful Stressful Neutral Not stressful Not very stressful	17	32.1%	36	67.9%	53	21.2%	18.093 (0.001)
		34	33.3%	68	66.7%	102	40.8%	
		16	47.1%	18	52.9%	34	13.6%	
		33	62.3%	20	37.7%	53	21.2%	
		6	75.0%	2	25.0%	8	3.2%	

Table 2 Test of Association Between Respondents' Awareness About the Symptoms of the Novel Coronavirus and Government Intervention Effectiveness in Dire Dawa City in the Year 2020

Symptoms of the Disease	Respondents Response Categories	Government Intervention Effectiveness to Cease COVID-19 in DDC						Chi-Square (Sig)
		Not Effective		Effective		Total		
		Count	%	Count	%	Count	%	
Fever	Not Related	7	70.0%	3	30.0%	10	4.0%	15.193 (0.001)
	Yes related	99	44.2%	125	55.8%	224	89.6%	
	I do not know	0	0.0%	16	100.0%	16	6.4%	
Cough	Not Related	1	25.0%	3	75.0%	4	1.6%	8.280 (0.016)
	Yes Related	105	44.5%	131	55.5%	236	94.4%	
	I do not know	0	0.0%	10	100.0%	10	4.0%	
Shortness of breath	Not related	2	33.3%	4	66.7%	6	2.4%	6.291 (0.043)
	Yes related	94	46.1%	110	53.9%	204	81.6%	
	I do not know	10	25.0%	30	75.0%	40	16.0%	
Sore throat	Not Related	10	76.9%	3	23.1%	13	5.2%	7.136 (0.028)
	Yes related	64	42.1%	88	57.9%	152	60.8%	
	I do not know	32	37.6%	53	62.4%	85	34.0%	
Headaches	Not Related	9	69.2%	4	30.8%	13	5.2%	11.438 (0.003)
	Yes related	94	43.7%	121	56.3%	215	86.0%	
	I do not know	3	13.6%	19	86.4%	22	8.8%	
Fatigue (tiredness)	Not Related	17	41.5%	24	58.5%	41	16.4%	19.651 (0.000)
	Yes related	85	49.7%	86	50.3%	171	68.4%	
	I do not know	4	10.5%	34	89.5%	38	15.2%	
Loss of taste and smell	Not Related	26	68.4%	12	31.6%	38	15.2%	14.495 (0.001)
	Yes related	53	41.7%	74	58.3%	127	50.8%	
	I do not know	27	31.8%	58	68.2%	85	34.0%	

Table 3 Test of Association Between Prevention Measures Practice Taken by the Respondents to Prevent Infection from the Novel Coronavirus and Government Intervention Effectiveness in Dire Dawa City in the Year 2020

Preventive Measures Taken by the Respondent to Prevent Infection from the Disease	Response Categories	Government Intervention Effectiveness to Cease COVID-19 in DDC						Chi-Square (Sig)
		Not Effective		Effective		Total		
		Count	%	Count	%	Count	%	
Hand washing for at least 20 seconds	No I do not Yes I do	27 79	77.1% 36.7%	8 136	22.9% 63.3%	35 215	14.0% 86.0%	20.115 (0.000)
Avoiding touching ones' body with unwashed hands	No I do not Yes I do	6 100	14.0% 48.3%	37 107	86.0% 51.7%	43 207	17.2% 82.8%	17.207 (0.000)
Use of disinfectant/Sanitizers to clean hands	No I do not Yes I do	27 79	33.8% 46.5%	53 91	66.2% 53.5%	80 170	32.0% 68.0%	3.604 (0.05)
Physical distancing (keeping minimum 2 meters or 6 feet)	No I do not Yes I do	16 90	61.5% 40.2%	10 134	38.5% 59.8%	26 224	10.4% 89.6%	4.352 (0.037)
Disinfecting surfaces	No I do not Yes I do	43 63	64.2% 34.4%	24 120	35.8% 65.6%	67 183	26.8% 73.2%	17.777 (0.000)

Additionally, the proportions of educators who rate the government intervention as effective corresponding cough as a symptom of the disease categories of not related and yes related were 75% and 55.5%, respectively. This signals that the proportions of respondent rate of government intervention effectiveness in scrubbing the disease less for who responded as yes it's among the symptoms of the pandemic than those who did not respond as it's among the symptoms of novel coronavirus (see Table 2).

Generally, eighty-four indicator variables were significantly associated with the government intervention effectiveness to cease COVID-19 (see Tables 1–12 respectively).

Multiple Logistic Regressions Analysis

By using multiple logistic regression (Forward Stepwise method), nine predictor variables were found to be significant: Not touching ones, organ with unwashed hands, disinfecting surfaces, fatigue (tiredness), getting the flu vaccine, an individual limitation in cooperating to cease the disease, how to maintain one's mental health during the isolation, hand washing practice for at least 20 seconds, family care practice of educators during the lockdown, and Self-isolation. Before going to the interpretation of the results, various tests were conducted to assess whether the specified model is well fitted to the observed data or not well fitted.

Assessing Logistic Regression Model

P-value < 0.001 (Table 13) shows that the model with predictor variables included has a better fit as compared to the model containing only a constant. This intern implies that the predictor variables were significantly affecting the outcome variable that was considered.

The Cox and Snell R-square indicate that 47.2% of the variation in the dependent variable is explained by the explanatory variables. Similarly, Nagelkerke's R-Square indicates that the 63.4% of the variability in the government's intervention measure effectiveness in ceasing the pandemic in Dire Dawa City was explained by the independent variables (see Table 14).

The test statistic has a chi-square value of 12.701 and a significance level of 0.123. This means that the Hosmer–Lemeshow test statistic is not statistically significant and, therefore, our model is quite a good fit to the data (see Table 15).

Out of the 250 educators included in the model 86.8% of them were correctly predicted which shows the overall percentage of prediction accuracy. The sensitivity is given by 84.7% and the specificity is given by 89.6%, which indicate that 84.7% % of the educators who did say the government intervention were effective and 89.6% of those who did not were correctly predicted in their respective categories.

Table 4 Test of Association Between Effective Preventive Methods of the Pandemic and Government Intervention Effectiveness in Dire Dawa City in 2020

Effective Preventive Methods of the Pandemic	Response Categories	Not Effective		Effective		Chi-Square (Sig.)
		Count	%	Count	%	
Being at home if there is sickness	No its not	6	10.9%	49	89.1%	28.633 (0.000)
	Yes its	99	51.3%	94	48.7%	
	I do not know	1	50.0%	1	50.0%	
Herbal supplements	No its not	57	43.5%	74	56.5%	7.084 (0.029)
	Yes its	30	34.1%	58	65.9%	
	I do not know	19	61.3%	12	38.7%	
Getting the flu vaccine	No its not	60	58.8%	42	41.2%	32.940 (0.000)
	Yes its	26	22.8%	88	77.2%	
	I do not know	20	58.8%	14	41.2%	
Using antibiotics	No its not	6	42.9%	8	57.1%	7.819 (0.020)
	Yes its	95	45.7%	113	54.3%	
	I do not know	5	17.9%	23	82.1%	
Using homeopathic remedies	No its not	18	29.5%	43	70.5%	6.978 (0.031)
	Yes its	23	54.8%	19	45.2%	
	I do not know	65	44.2%	82	55.8%	
Self-isolation	No its not	8	61.5%	5	38.5%	19.252 (0.000)
	Yes its	95	46.8%	108	53.2%	
	I do not know	3	8.8%	31	91.2%	
Disinfecting surfaces	No its not	10	71.4%	4	28.6%	5.316 (0.070)
	Yes its	81	41.3%	115	58.7%	
	I do not know	15	37.5%	25	62.5%	
Disinfecting the mobile phone	No its not	3	23.1%	10	76.9%	7.711 (0.021)
	Yes its	92	46.9%	104	53.1%	
	I do not know	11	26.8%	30	73.2%	

Table 5 Test of Association Between Short-Long Term Effects of Government Intervention and Its Effectiveness in Dire Dawa City in the Year 2020

Short and Long Term Effects of Government Intervention	Response Categories	Government Intervention Effectiveness to Cease COVID-19 in DDC						Chi-Square (Sig.)
		Not Effective		Effective		Total		
		Count	%	Count	%	Count	%	
		Count	%	Count	%	Count	%	
Sufficiency of government intervention measures	Not sufficient Yes its sufficient I do not know	104 2	56.2% 3.1%	81 63	43.8% 96.9%	185 65	74.0% 26.0%	55.615 (0.000)
Individual effectiveness in using government the prevention measure	Not effective Yes I am effective	78 28	50.6% 29.2%	76 68	49.4% 70.8%	154 96	61.6% 38.4%	11.175 (0.001)
Advantage of government intervention in scrubbing the disease	Reduce prevalence	17	34.0%	33	66.0%	50	20.0%	11.990 (0.035)
	Reduce Death rate or increasing recovery rate	4	30.8%	9	69.2%	13	5.2%	
	Helping who need any support	5	33.3%	10	66.7%	15	6.0%	
	Raising awareness of the community	35	36.8%	60	63.2%	95	38.0%	
	All	41	58.6%	29	41.4%	70	28.0%	
Importance rate of the intervention	Others	4	57.1%	3	42.9%	7	2.8%	11.796 (0.019)
	Not very important	4	36.4%	7	63.6%	11	4.4%	
	Not important	4	57.1%	3	42.9%	7	2.8%	
	Neutral	2	22.2%	7	77.8%	9	3.6%	
	Important	60	53.1%	53	46.9%	113	45.2%	
Limitation of government intervention to cease the disease	Very Important	36	32.7%	74	67.3%	110	44.0%	7.523 (0.057)
	It's not early applicable	18	45.0%	22	55.0%	40	16.0%	
	It's not consider all the real situation of the city	67	41.9%	93	58.1%	160	64.0%	
	It does not has any limitation or its good	13	32.5%	27	67.5%	40	16.0%	
	I do not know	8	80.0%	2	20.0%	10	4.0%	
Limitation of the surrounding community in co-operating to cease the disease	Never follow government preventive measures at all	57	44.9%	70	55.1%	127	50.8%	15.263 (0.002)
	Follow government preventive measures a bit	42	47.7%	46	52.3%	88	35.2%	
	Others	6	50.0%	6	50.0%	12	4.8%	
	I do not know	1	4.3%	22	95.7%	23	9.2%	
	Not follow government preventive measures at all	9	34.6%	17	65.4%	26	10.4%	
Individual limitation in cooperating to cease the disease	Not follow government preventive measures at all	53	33.3%	106	66.7%	159	63.6%	23.249 (0.000)
	Not follow government preventive measures	12	63.2%	7	36.8%	19	7.6%	
	Others	32	69.6%	14	30.4%	46	18.4%	
	I do not know							

Respondents attitude towards the 14 days quarantine	Not good attitude at all	8	50.0%	8	50.0%	16	6.4%	14.182 (0.007)
	Not good attitude	3	60.0%	3	40.0%	5	2.0%	
	Neutral	8	47.1%	8	52.9%	17	6.8%	
	Good attitude	74	48.7%	78	51.3%	152	60.8%	
	Very good attitude	13	21.7%	47	78.3%	60	24.0%	
Responses by surrounding community to help each other	Cloth Sharing	3	30.0%	7	70.0%	10	4.0%	13.728 (0.017)
	Meal sharing	34	46.6%	39	53.4%	73	29.2%	
	House rent exemption and postpone	4	44.4%	5	55.6%	9	3.6%	
	All	47	35.6%	85	64.4%	132	52.8%	
	I did not see nothing	18	69.2%	8	30.8%	26	10.4%	

Multiple Logistic Regression Model Analysis Results

The hypothesis test for individuals (Wald test): $H_0: \beta_j=0$ or $H_a: \beta_j \neq 0, j=1, 2, 3, \dots, k$.

As can be seen from Table 17, not touching ones with unwashed hands is significant at the 1% level. The value of $\exp(\hat{\beta}) = 11.940$ implies that the odds of government effectiveness are 11.940 times more likely for those respondents who did not refrain from touching one's body with unwashed hands as compared to the survey respondents who did avoid touching their eyes, nose, and mouth with unwashed hands held constant by the other predictors in the model.

For the case of the disinfecting surfaces, the result displayed under Table 17 indicates that it is significant at the 1% level. The value of $\exp(\hat{\beta}) = 0.157$ implies that the odds of the government intervention measure were effective in scrubbing the pandemic decrease by a factor of 15.7 for those respondents as they disinfecting a surface. This indicates that survey respondents who were disinfecting surfaces were less likely to rate the government intervention measure was effective in scrubbing the pandemic as compared to those responded as they do not disinfecting a surface.

For the case of the fatigue (tiredness), the result displayed under Table 17 indicates that it is significant at the 1% level. The value of $\exp(\hat{\beta}) = 0.069$ implies that the odds of the government intervention measure were effective in scrubbing the pandemic decrease by a factor of 6.9 for those respondents as they were say yes tiredness is the symptom of novel coronavirus pandemic. This indicates that survey respondents who were responded yes tiredness is the symptom of COVID-19 were less likely to rate the government intervention measure was effective in scrubbing the pandemic as compared to those responded as it is not the symptoms of novel coronavirus.

Similarly, the value of $\exp(\hat{\beta}) = 0.235$ implies that the odds of the government intervention measure were effective in scrubbing the pandemic decrease by a factor of 23.5 for those respondents as they were do not know that fatigue is symptoms for coronavirus pandemic. This indicates that survey respondents who responded as they did not know the fatigue is the symptom of COVID-19 were less likely to rate the government intervention measure was effective in scrubbing the pandemic as compared to those responded as it is not the symptoms of novel coronavirus.

Table 6 Test of Association Between Government Intervention Effectiveness to Cease the Disease and Its Cons on the Respondents' Educational Activities in Dire Dawa City in the Year 2020

Negative Impact of Intervention Methods Held for Mitigating the Pandemic in the City	Response Categories	Not Effective		Effective		Chi-Square (Sig.)
		Count	%	Count	%	
Teachers are obligated to stay at home	No its not Yes its I do not know	29 77 0	53.7% 41.2% 0.0%	25 110 9	46.3% 58.8% 100.0%	9.565 (0.008)
Respondents' Fear of unemployment	No its not Yes its I do not know	34 72 0	34.3% 48.6% 0.0%	65 76 3	65.7% 51.4% 100.0%	7.206 (0.027)
Stress	No its not Yes its I do not know	41 63 2	59.4% 36.0% 33.3%	28 112 4	40.6% 64.0% 66.7%	11.322 (0.003)
Un able to conduct Professional activities	No its not Yes its I do not know	17 87 2	23.6% 50.9% 28.6%	55 84 5	76.4% 49.1% 71.4%	15.987 (0.000)
Cancellation of Training Schedules	No its not Yes its I do not know	31 68 7	68.9% 35.1% 63.6%	14 126 4	31.1% 64.9% 36.4%	19.249 (0.000)
Loss of motivation teach again (in the long ran)	No its not Yes its I do not know	34 70 2	61.8% 37.2% 28.6%	21 118 5	38.2% 62.8% 71.4%	11.094 (0.004)

Table 7 Test of Association Between Government Intervention Effectiveness to Cease the COVID-19 and Its Positive Impact on the Respondents' Activities

Positive Impact of Intervention Measures Taken by the Government to Mitigate the Pandemic	Response Categories	Not Effective		Effective		Chi-Square (Sig.)
		Count	%	Count	%	
Teachers are able to give Family care	No I did not Yes I do	26 80	70.3% 37.6%	11 133	29.7% 62.4%	14.168 (0.001)
Have got time for watching movies	No I did not Yes I do I do not know	27 76 3	31.4% 47.8% 60.0%	59 83 2	68.6% 52.2% 40.0%	6.796 (0.033)
Have got time for doing physical activity	No I did not Yes I do	33 73	28.2% 54.9%	84 60	71.8% 45.1%	18.473 (0.000)

Based on the results displayed in Table 17, getting the flu vaccine is significant at the 1% level. The value of $\exp(\hat{\beta}) = 1.121$ implies that educators who have responded as getting flu vaccine is effective preventive measure of the disease were 1.121 times more likely to respond the intervention is effective as compared to those who said not and the value of $\exp(\hat{\beta}) = 4.124$ implies that educators who have responded as they did not know were 4.124 times more likely to respond the intervention is effective as compared to those who said not.

Similarly, respondent type of information mostly needed related to how to maintain one's mental health during the isolation is significant at the 1% level. The value of $\exp(\hat{\beta}) = 4.728$ implies that the odds of responding as the intervention is effective increases by a factor of 4.728 for those respondent educators who responded as they mostly needed about how to maintain one's mental health during the isolation as compared to those who did not mostly need information about how to maintain one's mental health during the isolation.

Table 8 Test of Association Between Government Intervention Effectiveness to Cease the COVID-19 and the Respondents' Trust in the Sources of Information in Their Reporting About the Outbreak

Respondents' Trust in the Sources of Information	Response Categories	Not Effective		Effective		Chi-Square (Sig.)
		Count	%	Count	%	
Public television stations	No I do not trust	28	57.1%	21	42.9%	6.392 (0.041)
	Yes I trust	76	38.4%	122	61.6%	
	I do not know	2	66.7%	1	33.3%	
Conversations with colleagues	No I do not trust	31	58.5%	22	41.5%	7.714 (0.021)
	Yes I trust	73	38.6%	116	61.4%	
	I do not know	2	25.0%	6	75.0%	
Consultation with health workers	No I do not trust	18	66.7%	9	33.3%	8.182 (0.017)
	Yes I trust	85	39.0%	133	61.0%	
	I do not know	3	60.0%	2	40.0%	
Private radio stations	No I do not trust	46	63.0%	27	37.0%	20.474 (0.000)
	Yes I trust	52	32.1%	110	67.9%	
	I do not know	8	53.3%	7	46.7%	
Official government press releases	No I do not trust	24	40.0%	36	60.0%	8.276 (0.016)
	Yes I trust	74	40.9%	107	59.1%	
	I do not know	8	88.9%	1	11.1%	
Medical institutions press releases	No I do not trust	23	62.2%	14	37.8%	8.829 (0.012)
	Yes I trust	78	38.0%	127	62.0%	
	I do not know	5	62.5%	3	37.5%	
Opinion polls	No I do not trust	46	55.4%	37	44.6%	8.954 (0.011)
	Yes I trust	54	36.7%	93	63.3%	
	I do not know	6	30.0%	14	70.0%	

Table 9 Test of Association Between Government Intervention Effectiveness to Cease the COVID-19 and Types Information Respondents Mostly Need Related to

Types Information Respondents Mostly Need Related to	Response Categories	Not Effective		Effective		Chi-Square (Sig.)
		Count	%	Count	%	
Personal stories from other people on how they cope with the pandemic situation	No Yes	21 85	26.2% 50.0%	59 85	73.8% 50.0%	12.564 (0.000)
How they can take care of a person who belongs to a risk group	No Yes	38 68	29.7% 55.7%	90 54	70.3% 44.3%	17.357 (0.000)
How they can best take care of their children's school education	No Yes	29 77	63.0% 37.7%	17 127	37.0% 62.3%	9.837 (0.002)
How the novel coronavirus is different from other diseases such as flu	No Yes	36 70	35.0% 47.6%	67 77	65.0% 52.4%	3.979 (0.046)
How they will be impacted economically by the pandemic	No Yes	15 91	19.0% 53.2%	64 80	81.0% 46.8%	25.923 (0.000)
How to maintain their mental health during the isolation time	No Yes	44 62	33.8% 51.7%	86 58	66.2% 48.3%	8.114 (0.004)

Table 10 Test of Association Between Government Intervention Effectiveness to Cease the COVID-19 and the Frequency of Respondent to Inform Themselves About the Novel Coronavirus per Day

Indicator Variable		Response Categories	Not Effective		Effective		Chi-Square (Sig.)
			Count	%	Count	%	
Frequency of Information respondents inform to themselves per day	Never	8	19.5%	33	80.5%		16.497 (0.002)
	Not Several times a day	8	53.3%	7	46.7%		
	Several times a day	33	37.9%	54	62.1%		
	Many Times a day	15	62.5%	9	37.5%		

From the results of logistic regression analysis, we can see that family care is a significant predictor of intervention effectiveness.

From the results, as an individual limitation in cooperating to cease the disease, we can see that respondents who responded not follow government preventive measures are 14.270 times more likely to rate the government intervention measures are effective as compared to those who responded as they did not follow government intervention measures at all. Similarly, respondents who responded others are 3.766 times more likely to rate the government intervention effectively as compared to those who responded as they did not follow government intervention measures at all, kept constant all other explanatory variables in the specified model.

Discussion

From the result of descriptive statistics analysis, 57.6% of the respondents were reported the intervention action applied by the government was effective to scrub COVID-19 in Dire Dawa City.

Based on the findings of bivariate analysis, out of indicator variables those have used in the survey; eighty-four of them were found to have a significant association with government intervention measure effectiveness in halting the spread of novel coronavirus, which are summarized and presented as follows: for various indicator variables: respondents' level of knowledge about the pandemic, respondents' susceptibility to be infected with the novel coronavirus, illness severity due to the disease if happen, knowledge how to protect from coronavirus, avoiding an infection with the novel coronavirus in the current situation, knowledge and

Table 11 Test of Association Between Government Intervention Effectiveness to Cease the COVID-19 and Respondents Use of Sources of Information to Stay Informed About Novel Coronavirus

Respondents Use of Sources of Information to Stay Informed About Novel Coronavirus	Response Categories	Not Effective		Effective		Chi-Square (Sig.)
		Count	%	Count	%	
Daily or weekly newspapers	Never Rarely Sometimes Often Always	34 9 37 18 8	38.2% 21.4% 56.1% 45.0% 61.5%	55 33 29 22 5	61.8% 78.6% 43.9% 55.0% 38.5%	15.309 (0.004)
Conversations with family and friends	Never Rarely Sometimes Often Always	2 18 27 25 34	50.0% 50.0% 30.3% 37.3% 63.0%	2 18 62 42 20	50.0% 50.0% 69.7% 62.7% 37.0%	16.308 (0.003)
Conversation with colleagues	Never Rarely Sometimes Often Always	9 23 32 16 26	60.0% 41.8% 34.0% 30.8% 76.5%	6 32 62 36 8	40.0% 58.2% 66.0% 69.2% 23.5%	23.639 (0.000)
Private television stations	Never Rarely Sometimes Often Always	8 6 45 26 21	42.1% 22.2% 45.5% 35.6% 65.6%	11 21 54 47 11	57.9% 77.8% 54.5% 64.4% 34.4%	13.323 (0.000)
Websites or online news pages	Never Rarely Sometimes Often Always	13 16 35 19 23	100.0% 18.4% 45.5% 52.8% 62.2%	0 71 42 17 14	0.0% 81.6% 54.5% 47.2% 37.8%	45.993 (0.000)
Social media (eg Facebook, Twitter, YouTube, WhatsApp)	Never Rarely Sometimes Often Always	6 6 36 39 19	75.0% 37.5% 55.4% 47.0% 24.4%	2 10 29 44 59	25.0% 62.5% 44.6% 53.0% 75.6%	19.236 (0.001)
Private radio stations	Never Rarely Sometimes Often Always	40 25 11 19 11	43.0% 50.0% 20.8% 52.8% 61.1%	53 25 42 17 7	57.0% 50.0% 79.2% 47.2% 38.9%	15.532 (0.004)
Official, government press releases	Never Rarely Sometimes Often Always	9 23 19 33 22	42.9% 63.9% 28.8% 37.9% 55.0%	12 13 47 54 18	57.1% 36.1% 71.2% 62.1% 45.0%	15.128 (0.004)
Medical institutions press releases	Never Rarely Sometimes Often Always	11 18 18 28 31	57.9% 56.2% 27.7% 33.7% 60.8%	8 14 47 55 20	42.1% 43.8% 72.3% 66.3% 39.2%	19.748 (0.001)

(Continued)

Table 11 (Continued).

Respondents Use of Sources of Information to Stay Informed About Novel Coronavirus	Response Categories	Not Effective		Effective		Chi-Square (Sig.)
		Count	%	Count	%	
Opinion polls	Never	58	47.5%	64	52.5%	16.417 (0.003)
	Rarely	15	24.2%	47	75.8%	
	Sometimes	9	39.1%	14	60.9%	
	Often	11	44.0%	14	56.0%	
	Always	13	72.2%	5	27.8%	
Celebrities and social media influencers	Never	17	53.1%	15	46.9%	10.340 (0.035)
	Rarely	30	41.7%	42	58.3%	
	Sometimes	29	34.1%	56	65.9%	
	Often	16	39.0%	25	61.0%	
	Always	14	70.0%	6	30.0%	

self-assessed adherence to preventive measures, feeling of respondents about the closeness of COVID-19, newness of the outbreak, spread-ability of the pandemic, stressfulness of the disease, and frequency of respondent to inform themselves about the novel coronavirus per day.

For respondents' awareness about the symptoms of the novel coronavirus: fever, cough, shortness of breath, sore throat, headaches, fatigue, loss of taste and smell.

Prevention measure practice taken by the respondents to prevent infection from the novel coronavirus: hand washing for at least 20 seconds, refrain from touching one's body with unwashed hands, use of disinfectant/sanitizers to clean hands, physical distancing (keeping minimum 2 meters or 6 feet), and disinfecting surfaces as preventive measure taken by the respondent.

For effective actions applied for prevention the outbreak in the city: being home at the time of sick, herbal supplement, getting the flu vaccine, using antibiotics, using homeopathic remedies, self-isolation, disinfecting surfaces considered as effective preventive method by the respondent, and disinfecting the mobile phone.

For short-long term effects of government intervention: sufficiency of government intervention measures, individual effectiveness in using government the prevention measure, advantage of government intervention in scrubbing the disease, importance rate of the intervention, limitation of government intervention to cease the disease, limitation of the surrounding community in cooperating to cease the disease, individual limitation in cooperating to cease the disease, respondents' attitude towards the 14 days quarantine, and responses by surrounding community to help each other.

For the negative impact of government intervention on the respondents' activities: teachers are obligated to stay at home, unemployment, stress, unable to conduct professional activities, cancellation of training schedules, and loss of motivation teach again (in the long run).

For positive impact of government intervention on the respondents' activities: teachers are able to give family care, have time for watching movies, and have time for doing physical activity.

The respondents' trust in the sources of information in their reporting about the outbreak: public television stations, conversations with colleagues, consultation with health workers, private radio stations, official, government press releases, medical institutions press releases, and opinion polls.

For types information respondents mostly need related to: personal stories from other people on how they cope with the pandemic situation, how they can take care of a person who belongs to a risk group, how they can best take care of their children's school education, how the novel coronavirus is different from other diseases such as flu, how they will be impacted economically by the pandemic, and how to maintain their mental health during the isolation.

For respondents' use of sources of information to stay informed about novel coronavirus: daily or weekly newspapers, conversations with family and friends, conversation with colleagues, private television stations, websites or online news pages, social media (eg, Facebook, Twitter, YouTube, WhatsApp), private radio station, official, government press releases, medical institutions press releases, opinion polls, and celebrities and social media influencers.

Table 12 Test of Association Between Government Intervention Effectiveness to Cease the COVID-19 and Trust (Perceptions) of Respondents in Institutions Those Inform Them About the Disease

Trust (Perceptions) of Respondents in Institutions Those Inform Them About the Disease	Response Categories	Not Effective		Effective		Chi-Square (Sig.)
		Count	%	Count	%	
Respondents Employer (if applicable)	Very Low Low Neutral High Very High	49 11 26 16 4	47.1% 22.9% 53.1% 39.0% 50.0%	55 37 23 25 4	52.9% 77.1% 46.9% 61.0% 50.0%	11.068 (0.026)
Media	Very Low Low Neutral High Very High	10 14 20 52 10	21.3% 58.3% 44.4% 43.7% 66.7%	37 10 25 67 5	78.7% 41.7% 55.6% 56.3% 33.3%	14.858 (0.005)
Other opinion leaders	Very Low Low Neutral High Very High	37 26 12 27 4	39.8% 34.2% 36.4% 67.5% 50.0%	56 50 21 13 4	60.2% 65.8% 63.6% 32.5% 50.0%	13.348 (0.010)
Ministry of Health	Very Low Low Neutral High Very High	3 8 4 54 37	33.3% 61.5% 25.0% 69.2% 27.6%	6 5 12 24 97	66.7% 38.5% 75.0% 30.8% 72.4%	39.227 (0.000)
Medical professional associations	Very Low Low Neutral High Very High	6 14 34 41 11	50.0% 35.9% 32.7% 56.2% 50.0%	6 25 70 32 11	50.0% 64.1% 67.3% 43.8% 50.0%	11.155 (0.025)
Schools	Very Low Low Neutral High Very High	15 27 26 29 9	39.5% 30.3% 51.0% 49.2% 69.2%	23 62 25 30 4	60.5% 69.7% 49.0% 50.8% 30.8%	11.907 (0.018)
Kindergartens	Very Low Low Neutral High Very High	40 17 13 30 6	38.5% 30.4% 40.6% 68.2% 42.9%	64 39 19 14 8	61.5% 69.6% 59.4% 31.8% 57.1%	16.004 (0.003)
Public transportation companies	Very Low Low Neutral High Very High	11 46 15 25 9	24.4% 46.0% 50.0% 54.3% 31.0%	34 54 15 21 20	75.6% 54.0% 50.0% 45.7% 69.0%	11.403 (0.022)
Police	Very Low Low Neutral High Very High	5 14 41 28 18	14.3% 32.6% 40.6% 62.2% 69.2%	30 29 60 17 8	85.7% 67.4% 59.4% 37.8% 30.8%	28.072 (0.000)

(Continued)

Table 12 (Continued).

Trust (Perceptions) of Respondents in Institutions Those Inform Them About the Disease	Response Categories	Not Effective		Effective		Chi-Square (Sig.)
		Count	%	Count	%	
Army	Very Low	14	63.6%	8	36.4%	25.814 (0.000)
	Low	37	37.4%	62	62.6%	
	Neutral	10	31.2%	22	68.8%	
	High	35	66.0%	18	34.0%	
	Very High	10	22.7%	34	77.3%	
The President/Prime Minister	Very Low	7	58.3%	5	41.7%	10.525 (0.032)
	Low	4	44.4%	5	55.6%	
	Neutral	26	59.1%	18	40.9%	
	High	46	41.8%	64	58.2%	
	Very High	23	30.7%	52	69.3%	

Table 13 Summary Statistics of the Likelihood Ratio Test

-2 Log Likelihood		Likelihood Ratio Tests.		
Null Model (Intercept Only)	Final Model	Chi-Square	D.F.	Sig.
340.775	159.485	181.290	14	0.000

Table 14 Model Summary

-2 Log Likelihood	Cox & Snell R Square	Nagelkerke R Square
181.290	0.472	0.634

Table 15 Hosmer and Lemeshow Test

Chi-square	D.F.	Sig.
12.701	8	0.123

For trust (perceptions) of respondents in institutions, those who inform them about the disease: respondents' employer (if applicable), media, other opinion leaders,

Ministry of Health, medical professional associations, schools, kindergartens, public transportation companies, police, army, and the president/prime minister.

In classical approach of binary logistic regression model only nine predictor variables were significant: refrain from touching one's body with unwashed hands, disinfecting surfaces, fatigue (tiredness), getting the flu vaccine, an individual limitation in cooperating to cease the disease, how to maintain one's mental health during the isolation, hand washing for at least 20 seconds, family care, and self-isolation (see Table 17).

Our study agrees with recent empirical studies on Humanitarian Impact, established multi-sectorial coordination meetings disrupted⁶ and with studies carried out in

Table 16 Classification Table for the Logistic Regression

Observed		Predicted		
		Government Intervention Effectiveness		Percentage
		Not Effective	Effective	
Government intervention effectiveness	Not effective	95	11	89.6
	Effective	22	122	84.7
Overall Percentage				86.8

Table 17 Variables in the Final Multiple Classical Binary Logistic Regression Model

Predictors	β	S.E. (β)	Wald	D.F.	Sig.	$\exp(\beta)$	95% C.I. for $\exp(\beta)$	
							Lower	Upper
Avoiding touching ones' body with unwashed hands (1)	2.5	0.755	10.803**	1	0.001	11.940	2.721	52.391
Disinfecting surfaces (1)	-1.9	0.505	13.441**	1	0.000	0.157	0.058	0.423
Fatigue (tiredness)			9.435**	2	0.009			
Fatigue (1)	-2.7	0.879	9.296**	1	0.002	0.069	0.012	0.384
Fatigue (2)	-1.5	0.707	4.192*	1	0.041	0.235	0.059	0.940
Getting the flu vaccine as an effective preventive method			9.919**	2	0.007			
Getting the flu vaccine (1)	0.11	0.562	0.041	1	0.839	1.121	0.373	3.372
Getting the flu vaccine (2)	1.42	0.586	5.850*	1	0.016	4.124	1.308	12.999
An individual limitation in cooperating to cease the disease			14.227**	3	0.003			
An individual limitation in cooperating to cease the disease (1)	2.66	0.736	13.051**	1	0.000	14.270	3.374	60.361
An individual limitation in cooperating to cease the disease (2)	1.33	0.506	6.866**	1	0.009	3.766	1.397	10.152
An individual limitation in cooperating to cease the disease (3)	0.99	0.683	2.103	1	0.147	2.692	0.706	10.265
How to maintain ones mental health during the isolation as mostly needed types of information (1)	1.55	0.500	9.657**	1	0.002	4.728	1.775	12.594
Hand washing practice for at least 20 seconds (1)	3.28	0.844	15.104**	1	0.000	26.628	5.088	139.367
Family care practice of educators during the lockdown as Pros of the government intervention	1.88	0.604	9.659**	1	0.002	6.544	2.001	21.396
Self-isolation as effective preventive method			5.733	2	0.057			
Self-isolation (1)	-2.2	1.165	3.708	1	0.054	0.106	0.011	1.041
Self-isolation (2)	-1.9	0.829	5.574*	1	0.018	0.141	0.028	0.717
Constant	-3.3	1.546	4.553*	1	0.033	0.037		

Notes: **Significant at the 1% level, *significant at the 5% level.




Figure 1 Bar chart, the respondent educators' level of knowledge about the pandemic in Dire Dawa City in 2020. From Figure 1, one can see that majority of the respondents have good knowledge about the pandemic in Dire Dawa City at the time of the study.

WHO region countries the delay in initiating travel restrictions, social distancing measures or lockdown and violations of measures lead to rapid the increase of cases and deaths in Iran.¹⁸

The following eight lessons were taken as important measures in reducing and controlling the disease:¹⁹ 1. acting early: as the pandemic is starting as small outbreaks but intensifying exponentially it is necessary to initiate action when the threat appears small. Decisions on reporting, travel restrictions, ban on mass/religious gatherings should be taken early, eg, KSA, Germany.¹⁹ 2. Epidemiological investigations, contact tracing and containment measures: countries with extensive testing and contact tracing like China, Singapore, South Korea and Germany showed better outcomes than countries with

limited testing like Italy.¹⁹ 3. Hospital preparedness: enhancing testing capacity, manpower training and increase in hospital resources like isolation wards, ICU and ventilators as done by countries like Germany, which have a low CFR, Singapore and China.¹⁹ 4. Monitoring and reporting: essential to have a data documentation and dissemination process to plan the resources wisely and provide right information to people, eg, Singapore, China.¹⁹ 5. Low-cost innovations in patient testing and treatment, eg, Sri Lanka.¹⁹ 6. Safety of healthcare staff: not testing the healthcare staff on priority and shortage of PPE led to nosocomial infections and deaths among doctors as in Italy.¹⁹ 7. Stringent social distancing measures: total lockdown has better outcomes than phase-wise lockdown as observed in KSA in comparison with Iran.¹⁹ 8. A




Figure 2 The rating of the timing of government intervention measures to scrub the disease in Dire Dawa City. From Figure 2, one can visualize that 49.2% of the respondent educators rate the government's intervention measures timing was not early in Dire Dawa City.

prior experience of dealing with pandemics enables better preparedness and outcomes, eg, Singapore and KSA.¹⁹

Additionally, a study done in Italy by Giordano et al²⁰ demonstrated that restrictive social-distancing measures will need to be combined with widespread testing and contact tracing to end the ongoing COVID-19 pandemic²⁰ which is somehow in accordance with our study findings.

As another study done by Sweden ECDC²¹ implied that the preexisting number of hospital beds/1000 population in Germany was 8.3 when compared to 3.4 in Italy.²¹

Additionally, our study finding, among the types of information respondents need mostly related to, how to maintain one's mental health during the isolation time had an effect on the government intervention effectiveness, which can be taken in line with the results of the study conducted by Alvarez-Risco et al¹¹ showed that Peru's healthcare workers' anxiety and mental distress decreased as the distance from the epicenter increases, corroborating the ripple effect and disconfirming the

typhoon eye theory.¹¹ A lower education level increased the anxiety levels, whereas age and gender did not affect the anxiety and distress levels.¹¹ Alvarez-Risco et al¹¹ have reported that their results can help guide mental health service providers toward vulnerable groups of healthcare workers that are closer to Lima; the COVID-19 epicenter in Peru.¹¹

Similarly, anxiety of the educators was associated with government effectiveness in our study. Thus, our finding was in accordance with the study done by Alvarez-Risco et al.¹⁰

Further, the study made by Alvarez-Risco et al¹² on the COVID-19 pandemic created an opportunity to expand telemedicine services. Alvarez-Risco et al¹² foresee that telemedicine in Peru could help fortify disease prevention programs, monitor chronic disease patients, and combat malnutrition in vulnerable populations, such as pregnant women and children under 5 years old. Peru was forced into telemedicine due to the pandemic, and the government is working to improve Internet coverage.¹² This also




Figure 3 Bar chart shows the respondents, level of worry or feeling level about COVID-19 in Dire Dawa City. From Figure 3, one can visualize that 43.6% of the respondent educators' rate they felt worrying about the pandemic in Dire Dawa City.

resembles our study finding since postgraduate classes were ran via online platform, and course materials were uploaded online for the undergraduate university students where the government was increased the access of the Internet to the students and to the teachers. Where the difference in telemedicine and online education services is considered.

Another study done by Alvarez-Risco et al¹⁴ taking evidence from four countries: Colombia, Ecuador, Mexico, and Peru implied that COVID-19 has changed the world,¹⁴ in creating the need for new actions from society, including universities and companies¹⁴ which agrees with our study.

A study done by Yáñez et al¹⁵ implied that self-medication affects both negatively and positively the health of people, which has become evident during the COVID-19 pandemic.¹⁵ Yáñez et al¹⁵ reported that there were

significant percentages of self-medication, including drugs without sufficient scientific evidence, age, region where one lived and job status was variables associated with self-medication frequency.

Finally, Yáñez et al¹⁵ suggested that a continuous awareness and sensitization about the risks of self-medication are warranted which have more similar finding with our study in terms of respondents' perception to the symptoms, preventive methods, need of awareness creation and effective preventive methods of the pandemic. where it differs in terms of demographic variables since demographic variables were not included in our questionnaires adopted from WHO standard questionnaires.

Similarly, the study performed by Yan et al¹⁶ signaled that the COVID-19 outbreak has highlighted the need to target infodemics that can be as detrimental as an actual

epidemic. It will be a multifactorial fight because they will need to increase health literacy in the population, establish a stronger presence of national health agencies in social media, develop better detection tools, and enable action by governments, as Peru has implemented. This finding is also similar to ours because knowledge and perceptions of the respondents were significantly associated with government intervention measure effectiveness.

Further, as study done by Alvarez-Risco et al¹⁷ implied Godlee appraises the UK government's response to COVID-19. On 18 May the Peruvian government started allowing supervised walks for people under 14 years old. Alvarez-Risco et al¹⁷ have previously expressed their concern for this measure, because it could lead to a rise in cases in children, who typically exhibit milder symptoms. Alvarez-Risco et al¹⁷ reported that there must be cautious in Peru, where there is a precarious health system, cases keep increasing in small children, paediatric hospitals are saturated with adult patients, and public policies are not necessarily following the global epidemiological alerts. This finding is in agreement with our study finding in terms of the need of cautious in Dire Dawa City too at the time of the study.

A study by Yan et al¹⁶ signaled that the hospitality industry worldwide is suffering under the COVID-19 pandemic which is in parallel with our study finding that education institutes were suffering too under the pandemic in the Dire Dawa City.

On the other hand, a paper done by UNOCHA⁶ showed that

the impacts of education development, conceptual development, and country supports on the entrepreneur's ability to carry out green entrepreneurship were positive.⁶ The research findings also may be helpful for the governments in establishing new norms to promote entrepreneurship which is in line with our study that government can use new strategy to alleviate the pandemic.

Conclusion

Based on descriptive analysis of indicator 57.6% of the study participants reported as the government was effective in halting the pandemic in the city. On the other side, from bivariate analysis result, eighty-four indicator variables have a significant association with government's intervention effectiveness in halting the pandemic.

On the bases classical binary logistic regression analysis nine predictors significantly related with the government's intervention effectiveness in halting the pandemic

in the study area: refrain from touching one's body with unwashed hands, disinfecting surfaces, fatigue (tiredness), getting the flu vaccine, individual limitation in cooperating to cease the disease, how to maintain one's mental health during the isolation, hand washing practice of respondents' for at least 20 seconds, respondents got time for family care during lockdown, and self-isolation were factors most probable affects government intervention effectiveness as shown in the discussion part of the document. We recommend that every concerned body has to give a considerable attention to prevention method of the disease in the study area.

Dire Dawa City Administration and the concerned body have to give the appropriate attention to the awareness of teachers about symptoms of the pandemic; to obtain an expected change.

Similarly, respondents' individual limitation in cooperating with government intervention measures to cease the disease has an important impact on the government intervention effectiveness. Therefore, it is better to improve the cooperating habit of the educators to the preventive measures of novel coronavirus in Dire Dawa City since individual has Lion's share in mitigating the spread of the outbreak.

Additionally, among the types of information respondents need mostly related to, how to maintain one's mental health during the isolation time had an effect on the government intervention effectiveness. Thus, appropriate and timely measures need to be taken by the Ministry of Health and other concerned bodies in providing advices and counseling regarding types of information respondents need mostly related about during the isolation time period.

Eventually, regarding the advantage of the outbreak the following indicators were significant: respondents were able to get time for reading, respondents were able to get chance of Family care practice, respondents were able to get chance of watching movies, and respondents were able to get involved in physical exercise activates were among the importance brought by the pandemic to the educators due to the lockdown intervention actions applied in the city.

On the contrary, the following indicators have significant negative impact on educational activities of respondents: teachers are obligated to stay at home, fear of unemployment, stress, unable to conduct professional activities, cancellation of training, and loss of motivation.

Thus, the concerned bodies in the city have to work together to alleviate such problems and to have a well and

productive teacher by considering the intensity of the negative impacts of the pandemic in the city since it has both short- and long-term negative impacts on respondents' success. Generally, the cumulative success of the study participants can be considered as the government intervention success in study area.

Abbreviations

COVID-19, Coronavirus Disease 2019; JHU, Johns Hopkins University; MERS, Middle East Respiratory Syndrome; SARS, Severe Acute Respiratory Syndrome; UNOCHA, United Nations Office for the Coordination of Humanitarian Affairs; WHO, World Health Organization; WB, World Bank; and AGDH, Australian Government Department of Health.

Acknowledgments

The authors would like to thank Dire Dawa University: Department of Statistics, College of Natural and Computational Sciences, College of Social Sciences and Humanities, College of Medicine and Health Sciences, Research Affairs Directorate, Community Service and Research Affairs Vice President.

Additionally, we would like to forward our appreciation to Dawa Administration Health Bureau, Dire Dawa Administration Education Bureau, School Principals, School Coordinators, and School owners for their warm welcoming and support during data collection stage.

Finally, we are highly grateful for all participants, enumerators and data feeders.

Author Contributions

All authors participated in data analysis, preparing, reconsidering, and finalizing the article, gave final approval of the version to be published, agreed to the submitted journal, and agreed to be fully accountable for all aspects of the work.

Disclosure

The authors declare that they have no conflicts of interest in this study.

References

1. Australian Government Department of Health. COVID-19 – Frequently asked questions; 2020. Available from: <https://www.bluecross.com.au/mvc/k12webapi/media/pdf/covid-19/coronavirus-covid-19-frequently-asked-questions.pdf>. Accessed September 15, 2021.
2. WHO. Global situation daily report; 2021. Available from: <https://www.who.int/>. Accessed September 3, 2021.
3. DDA. Dire-Dawa-administration-health-bureau daily COVID-19 situational report; 2021. Available from: <https://www.facebook.com/>. Accessed September 3, 2021.
4. JHU. Daily report of global coronavirus pandemic prevalence; 2020. Available from: <https://coronavirus.jhu.edu/>. Accessed September 3, 2021.
5. Liya T. Ethiopian ministry of health novel coronavirus daily report; 2020. Available from: <https://www.moh.gov.et/ejcc/index.php/en>. Accessed September 3, 2021.
6. UNOCHA. ETHIOPIA: COVID-19 humanitarian impact situation update No. 01; 2020. Available from: <https://www.humanitarianresponse.info/en/operations/ethiopia/covid-19>. Accessed September 3, 2021.
7. Zhang SX, Chen J, Jahanshahi AA, et al. Succumbing to the COVID-19 pandemic-healthcare workers not satisfied and intend to leave their jobs. *Int J Ment Health Addict*. 2021;1–10. doi:10.1007/s11469-020-00418-6
8. Zhang SX, Sun S, Jahanshahi AA, et al. Developing and testing a measure of COVID-19 organizational support of healthcare workers – results from Peru, Ecuador, and Bolivia. *Psychiatry Res*. 2020;291:113174. doi:10.1016/j.psychres.2020.113174
9. Yáñez JA, Jahanshahi AA, Alvarez-Risco A, et al. Anxiety, distress, and turnover intention of healthcare workers in Peru by their distance to the epicenter during the COVID-19 crisis. *Am J Trop Med Hyg*. 2020;103(4):1614. doi:10.4269/ajtmh.20-0800
10. Alvarez-Risco A, Del-Aguila-Arcentales S, Yáñez JA. Perspective piece telemedicine in Peru as a result of the COVID-19 pandemic: perspective from a country with limited internet access. *Am J Trop Med Hyg*. 2021;105(1):6–11. doi:10.4269/ajtmh.21-0255
11. Alvarez-Risco A, Estrada-Merino A, de Las Mercedes Anderson-seminario M, et al. Multitasking behavior in online classrooms and academic performance: case of university students in Ecuador during COVID-19 outbreak. *Interact Technol Smart Educ*. 2020. Available from: <https://www.emerald.com/insight/publication/issn/1741-5659>. Accessed September 3, 2021.
12. Alvarez-Risco A, Del-Aguila-Arcentales S, Rosen MA, et al. Expectations and interests of university students in COVID-19 times about sustainable development goals: evidence from Colombia, Ecuador, Mexico, and Peru. *Sustainability*. 2021;13(6):3306. doi:10.3390/su13063306
13. Aleidi S, Issa A, Bustanji H, Khalil M, Bustanji Y. Self-medication practices during the COVID-19 pandemic among the adult population in Peru: a cross-sectional survey. *Saudi Pharm J*. 2015;23(3):250–256. doi:10.1016/j.jsps.825
14. Alvarez-Risco A, Mejia CR, Delgado-Zegarra J, et al. The Peru approach against the COVID-19 Infodemic: insights and strategies. *Am J Trop Med Hyg*. 2020;103(2):583–586. doi:10.4269/ajtmh.20-0536
15. Yáñez JA, Alvarez-Risco A, Delgado-Zegarra J. Covid-19 in Peru: from supervised walks for children to the first case of Kawasaki-like syndrome. *BMJ*. 2020;369. doi:10.1136/bmj.m2418
16. Yan J, Kim S, Zhang SX, et al. Hospitality workers' COVID-19 risk perception and depression: a contingent model based on transactional theory of stress model. *Int J Hosp Manag*. 2021;95:102935. doi:10.1016/j.ijhm.2021.102935
17. Alvarez-Risco A, Młodzianowska S, García-Ibarra V, et al. Factors affecting green entrepreneurship intentions in Business University Students in COVID-19 pandemic times: case of Ecuador. *Sustainability*. 2021;13(11):6447. doi:10.3390/su13116447
18. WHO. Coronavirus disease situation reports; 2020. Available from: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports>. Accessed September 3, 2021.
19. Davalgi S, Undi M, Annadani R, et al. Comparison of measures adopted to combat COVID 19 pandemic by different countries in WHO regions. *Indian J Community Health*. 2020;32:288–299.

20. Giordano G, Blanchini F, Bruno R, et al. Modeling the COVID-19 epidemic and implementation of population-wide interventions in Italy. *Nat Med*. 2020;26(6). doi:10.1038/s41591-020-0883-7
21. Sweden ECDC. Updated rapid risk assessment from ECDC on the novel Coronavirus disease 2019 (COVID-19) pandemic: increased transmission in the EU/EEA and the UK, Stockholm. *Eurosurveillance*. 2020;25(10):2003121. doi:10.2807/1560-7917.ES.2020.25.12.2003261

Journal of Multidisciplinary Healthcare

Dovepress

Publish your work in this journal

The Journal of Multidisciplinary Healthcare is an international, peer-reviewed open-access journal that aims to represent and publish research in healthcare areas delivered by practitioners of different disciplines. This includes studies and reviews conducted by multidisciplinary teams as well as research which evaluates the results or conduct of such teams or healthcare processes in general. The journal

covers a very wide range of areas and welcomes submissions from practitioners at all levels, from all over the world. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-inflammation-research-journal>