

Assessment of the Critical Care Work Environment of Intensive Care Unit Nurses in Saudi Arabia

Mohammed Ageel , Abdullah Shbeer

College of Medicine, Jazan University, Jazan, Saudi Arabia

Correspondence: Mohammed Ageel, College of Medicine, Jazan University, Jazan, Saudi Arabia, Tel +966505769570, Email ageelahmed@jazanu.edu.sa

Background: Nurses play a major role in the delivery of complex and challenging critical care in intensive care units (ICUs). Assessment of work environment is essential indicators of hospital management and can be applied to workforce planning and identifying nursing profession needs. The American Association of Critical-Care Nurses (AACN) recognized six standards for a healthy work environment and developed the Healthy Work Environment Assessment Tool (HWEAT). The aim of this study was to assess the work environment of ICU nurses in Jazan, Saudi Arabia.

Methods: This cross-sectional study was conducted at public and private hospitals. Data were collected using a self-administered questionnaire that included the sociodemographic characteristics and the AACN HWEAT. Data were analyzed to obtain descriptive and inferential statistics. The Mann–Whitney U and Kruskal–Wallis tests were employed to compare demographic data, on the basis of the overall mean HWEAT score.

Results: The study participants were 238 ICU nurses, who were predominantly female (83%). The mean overall HWEAT score was 3.55 ± 1.03 , which is within the “good” range, and was higher for male nurses (3.66), nurses aged ≥ 41 years (3.76), and nurses with postgraduate education (4.04), work experiences of >10 years (3.63), and alternate work shifts (3.6). Nurses in private hospitals had significantly higher overall scores than nurses in public hospitals (3.83 vs 3.19, $P < 0.001$). All HWEAT standards (“effective decision-making”, “authentic leadership”, “appropriate staffing”, “true collaboration”, “skilled communication”, and “meaningful recognition”) were rated as good (mean range, 3.43–3.63).

Conclusion: The study results could assist hospitals in prioritizing the adoption of AACN HWE standards. A target benchmark of “good” was established for both the overall score and each standard, which indicates a good work environment as perceived by ICU nurses. In Addition, interpersonal differences should be considered when developing improvement initiatives.

Keywords: work environment, nursing practice, intensive care

Introduction

Nurses play a major role in the delivery of critical care in intensive care units (ICUs), where the care they provide is more complex and challenging than that provided in other hospital departments.¹ ICU nurses are frequently subjected to high demands to fulfill the duties assigned to them.² Working in a field that involves complex multitasking, high workloads, and providing specialized care to critically ill and reliant patients can be overwhelming for nurses,³ especially during the coronavirus disease (COVID-19) pandemic. The pandemic of COVID-19 has been the most disruptive factor in the ICU work environment. Over the past two years, ICU teams around the world have been inundated with critically ill patients requiring breathing support and other advanced interventions.⁴ It has been proven that the health of the work environments of critical care nurses affects patient care outcomes and the job satisfaction and retention of registered nurses.⁵ A growing body of evidence demonstrates a connection between nurse work environments and patient outcomes.^{6,7}

Work environment is generally defined as “the physical, chemical, biological, organizational, social, and cultural factors that surround a worker”.⁸ According to the American Association of Critical-Care Nurses (AACN), “a healthy

work environment (HWE) is imperative to ensure patient safety, enhance staff satisfaction and retention, and maintain an organization's financial viability".⁹ The ideal work environment provides conditions for physical, mental, and social well-being.¹⁰ The external pressures of diminishing health-care reimbursement, increasing regulatory requirements, rising acuity and patient complexity, and increased health-care operational restrictions have a negative effect on work environments.¹¹

Assessment of work environment is essential indicators of hospital management and can be applied to workforce planning and identifying nursing profession needs.¹² Clinical leaders have recognized that to maintain patient care quality, attention should be paid to the performance of health-care teams.⁷ The AACN has identified systemic behaviors that promote patient safety, optimal outcomes, and nursing practice excellence and classified them into six evidence-based standards that are critical for establishing and maintaining a HWE: "skilled communication", "true collaboration", "effective decision-making", "appropriate staffing", "meaningful recognition", and "authentic leadership".⁹ For this purpose, the AACN developed a reliable and valid tool, the Healthy Work Environment Assessment Tool (HWEAT), which comprehensively captures work environments to achieve and sustain environments of excellence.¹³

The impact of the nursing work environment is an important field of nursing research, especially because it is related to patient safety and quality of care.^{14–16} Nurses and health-care administrators must endeavor to improve the nursing work environment.¹⁷ Nursing research has progressively demonstrated the impact of the nursing work environment on the quality of nurses' work experiences and the delivery of quality patient outcomes in acute-care hospitals.^{18–20} However, research on the assessment of the work environment of ICU nurses after the COVID-19 pandemic is still scarce, especially in Saudi Arabia, which is needed to emphasize the patient safety and quality of care. Therefore, the aim of this study was to assess the work environment of nurses working in ICUs in Saudi Arabia using the HWEAT. The results of this study can provide baseline data to enable hospital management to benchmark their status based on established quality standards and may be useful for identifying HWE components that require improvement.

Methods

Population and Study Area

A cross-sectional study design was conducted at public and private hospitals affiliated with the Jazan Region Health Administration in Saudi Arabia, between January 2022 and June 2022. The survey population included full-time ICU nurses. Using G*Power software, the minimum sample size required to achieve 80% power for detecting a medium effect, at a significance criterion of $\alpha = 0.05$, was $N = 130$, for the Independent Sample *T*-Test.²¹ Owing to the lack of accessibility of the ICU nurses' contact information, nursing supervisors in the hospitals were contacted and asked to distribute the survey to ICU nurses in their hospitals. The questionnaire was distributed online using Qualtrics Research Core (Qualtrics, Provo, UT). The survey link was emailed to the nurse supervisors for them to forward to the nurses in their teams. A total of 326 invitations via email were sent to ICU nurses in 13 hospitals in January 2022, with a three follow-up emails every two months. Convenience samples of 238 were collected for this study, which is considered adequate.

Survey

Data were collected using a self-administered questionnaire that included the sociodemographic characteristics (age, sex, educational level, years of experience, time of shift, and type of hospital) and the AACN HWEAT to assess the work environment of the ICU nurses. The HWEAT is an 18-item Likert scale that ranks standard adherence as "needs improvement", "good", or "excellent." The 18-items evaluates each of the six HWE standards with three unique items. On a 5-point Likert scale, participants were requested to choose their level of agreement or disagreement with each statement (1 = "strongly disagree," 5 = "strongly agree").

The cover page of the survey questionnaire included an introduction sheet that contained the study's objective, duration, privacy and confidentiality statements. The first page of the questionnaire included the participation consent form, which participants should sign to access the survey.

Statistical Analyses

For the statistical analysis, Statistical Package for Social Sciences (SPSS) Version 27 (IBM Corp., Armonk, NY) was utilized to obtain the descriptive statistics for continuous variables (mean, standard deviation [SD]) and categorical variables (n, percentage). Responses from participants were averaged to get the mean score for each item, the mean score for each HWE standard, and the overall mean HWEAT score. The mean score was interpreted as categorical thresholds as follows: 4.00 to 5.00, “excellent”; 3.00 to 3.99, “good”; and 1.00 to 2.99, “needs improvement.”⁷ The percentage distributions of the ICU nurses at each HWEAT categorical threshold for the HWE standards were obtained. The Kolmogorov–Smirnov goodness-of-fit test is used for testing normality of the distributions of the quantitative variables. The Mann–Whitney U and Kruskal–Wallis tests were employed to compare demographic data. To evaluate statistical significance, a p value of 0.05 was applied.

Ethical Considerations

This study complies with the Declaration of Helsinki and was approved by the Ethics Committee of Jazan University (Approval No.: REC-43/11/279).

Results

This study included a total of 238 nurses working in ICUs, and the response rate was approximately 73%. As shown in Table 1, the sociodemographic statistics showed that 83% of the ICU nurses were female, 50% belonged to the age group of 22–30 years, 80% had a bachelor’s degree, 45% had 6–10 years of work experience, 65% worked alternate shifts, and 55% were employed in private hospitals. The mean overall HWEAT score was 3.55 ± 1.03 , which is within the “good” range. When the groups in each demographic category were compared, the overall HWEAT score was higher for male

Table 1 Summary of the Participants’ Demographic Characteristics and the Mean Overall HWEAT Score

Item		n (%)	Overall Mean (SD)	P value
Sex ¹	Female	198 (83)	3.52 (0.93)	0.448
	Male	40 (17)	3.66 (1.07)	
Age (years) ²	22–30	118 (50)	3.4 (1.08)	0.39
	31–40	104 (44)	3.68 (0.78)	
	≥41	16 (6.7)	3.76 (0.88)	
Qualification ²	Diploma	42 (18)	3.56 (0.96)	0.53
	Bachelor	190 (80)	3.53 (0.95)	
	Postgraduate	6 (2.5)	4.04 (0.91)	
Length of work experience (years) ²	0–5	83 (35)	3.47 (1.02)	0.82
	6–10	107 (45)	3.57 (0.90)	
	>10	48 (20)	3.63 (0.96)	
Typical work shift ²	Days	50 (21)	3.51 (1.04)	0.75
	Nights	34 (14)	3.35 (0.95)	
	Alternate	154 (65)	3.6 (0.93)	
Type of hospital ¹	Public	106 (45)	3.19 (1.01)	<0.001
	Private	132 (55)	3.83 (0.80)	

Note: ^aMann–Whitney U-test. ^bKruskal–Wallis test.

nurses (3.66), nurses aged ≥ 41 years (3.76), nurses with postgraduate education (4.04), nurses with work experience of >10 years (3.63), and nurses with alternate work shifts (3.6). ICU nurses in private hospitals had significantly higher overall scores than ICU nurses in public hospitals (3.83 vs 3.19, $P < 0.001$).

Table 2 shows the results for the responses to each item of the HWEAT. The item “Able to influence policies, procedures, and bureaucracy” scored highest (3.72). The items that scored lower than the mean overall HWEAT score were “Consistent use of data-driven, logical decision-making process”, “Enough staff to maintain patient safety”, “Formal recognition system makes staff feel valued”, and “Zero tolerance for disrespect and abuse” (3.17) in this order. All six standards scored within the “good” range, ranked from highest to lowest in terms of aggregate scores as

Table 2 Mean Scores of the Healthy Work Environment Assessment Tool Variables

Variable	Items	Mean (SD)
Skilled Communication	“Maintains frequent communication”	3.68 (1.02)
	“Matches actions to words”	3.55 (1.03)
	“Observes zero tolerance for disrespect and abuse”	3.17 (1.20)
	Overall	3.47 (0.92)
True Collaboration	“Involves staff in decision-making”	3.59 (1.04)
	“Influences policies, procedures, and bureaucracy”	3.72 (1.06)
	“Seeks input for decision-making”	3.58 (1.05)
	Overall	3.63 (1.04)
Effective Decision-making	“Consistent use of a data-driven, logical decision-making process”	3.48 (1.03)
	“Involves the right departments, professions, and groups”	3.67 (1.00)
	“Considers the patient’s perspective is in important decisions”	3.67 (0.96)
	Overall	3.61 (0.95)
Appropriate Staffing	“Enough staff to maintain patient safety”	3.30 (1.23)
	“Right mix of nurses and other staff to ensure optimal outcomes”	3.48 (1.13)
	“Support service level allows nurses and staff to focus on care”	3.51 (1.20)
	Overall	3.43 (1.11)
Meaningful Recognition	“Formal recognition system makes staff feel valued”	3.45 (1.09)
	“Staff members let people know when they have done a good job”	3.52 (1.12)
	“Motivating opportunities for personal growth”	3.56 (1.16)
	Overall	3.51 (1.08)
Authentic Leadership	“Staff have a positive relationship with nurse leaders”	3.63 (1.16)
	“Nurse leaders understand the dynamics at point of care”	3.60 (1.12)
	“Nurse leaders play a role in making key decisions”	3.67 (1.16)
	Overall	3.63 (1.11)




Figure 1 Percentage distributions of the nurses among the HWEAT categorical thresholds for the HWE standards.

follows: “true collaboration” and “authentic leadership” (3.63), “effective decision-making” (3.61), “meaningful recognition” (3.51), “skilled communication” (3.47), and “appropriate staffing” (3.43).

As shown in Figure 1, the percentage distribution of ICU nurses at each HWEAT categorical thresholds indicates that appropriate staffing standard was rated “needs improvement” by the highest percentage of nurses (21%), followed by meaningful recognition and skilled communications (18%). The item “effective decision-making” and “authentic leadership” “standards” was rated “excellent” by the highest percentage of ICU nurses (59%), followed by true “collaboration standards” (57%).

Discussion

Evidence has shown that work environments are strongly associated with patient safety, organizational financial stability, employment and retention.^{9,22} Assessment of work environments must be recognized as a main strategic and operational goal of health-care systems.¹³ In this study, the ICU nurses reported that the health of their work environment was “good”, rating each standard also as “good.” This result is consistent with those of earlier research.^{13,23,24} However, the mean scores reported in this study (range: 3.43–3.63) were lower than those reported (range: 3.53–3.76) in a previous study conducted in 2012 in Riyadh, Saudi Arabia.²⁴

The stratification of the responses according to demographic characteristic showed significantly more positive evaluations by male nurses (3.66), nurses aged ≥ 41 years (3.76), nurses with postgraduate education (4.04), nurses with a work experience of >10 years (3.63), and nurses with alternate work shifts (3.6). Nurses in private hospitals had significantly higher overall scores than nurses in public hospitals (3.83 vs 3.19, $P < 0.001$). Moreover, this study considered the percentage of nurses at each HWEAT categorical threshold, which showed that appropriate staffing standard was rated “needs improvement” by the highest percentage of nurses (21%). These may point to an opportunity for developing improvement plans that take into consideration interpersonal differences.²⁵

None of the HWE standards was rated “excellent.” The HWE is multifaceted, and a large body of research has shown that excellence in the work environment arises from a strong commitment to all six standards.⁹ Each HWE standard needs continuing attention and should be considered as an on-going improvement opportunity.⁹ Skilled communication is crucial in providing excellent patient care and job satisfaction.^{9,26}

Poor communication is a leading cause of burnout,^{27,28} and practitioners who get training in communication are less likely to have burnout.^{16,29,30} Despite that skilled communication was interpreted as good, the item “zero tolerance for disrespect and abuse” showed the lowest score (3.17) among all items in the HWEAT, which presents a need for further attention. According to Matthew Grissinger, “disrespectful behavior chills communication and collaboration, undercuts

individual contributions to care, undermines staff morale, increases staff resignations and absenteeism, creates an unhealthy or hostile work environment, causes some to abandon their profession, and ultimately harms patients”.^{30,31} Several studies have linked these behaviors to adverse events, medical errors, patient safety compromises, and even patient mortality.^{32,33} True collaboration is vital, with 90% of nurses believe that true collaboration is the most crucial aspect for a HWE; it can reduce compassion fatigue and result in improved patient care outcomes.^{29,34} In contrast, medical errors and staff turnover is linked to poor collaboration.³⁵ Effective decision-making enhances patient outcomes and compassion satisfaction while reducing compassion fatigue.^{36,37} It is recommended that employees be given chances to engage in decision-making processes, which should be matched with their corresponding accountability, in order to motivate them to strive for excellence.³⁸ Moreover, staff empowerment, specifically psychological empowerment, has significant impact of employees performance.³⁹ Appropriate staffing occurs when patient needs align with nurse competencies; mismatches increase the chance of complications and diminish nurses’ well-being.^{6,7} Nurses with high patient loads experience burnout.²⁰ A study found that 48% of nurses who quit their jobs cited poor or inadequate staffing as the reason.⁶ Moreover, studies have indicated that increasing the number of nurse staff assigned to a unit provides better patient care and decreases the incidence of mortality and adverse patient outcomes.^{20,40–42} Lack of meaningful recognition can be a source of compassion fatigue,⁹ while recognizing contributions³⁶ can result in compassion satisfaction.⁴³ A study indicated that providing personnel with opportunities to demonstrate pride in their work, to be recognized for their achievements, and to be encouraged by superiors strengthened the workforce, which eventually improved the work environment.⁴⁴ Authentic leadership is nursing leaders viewed as collaborators, team builders and skilled communicators, which their performance may impact nurse retention, job satisfaction and shaping the work environment.^{9,37,45}

The strengths of this study are that a reliable and valid assessment tool, the AACN HWEAT, was used to support continuous improvement through regular assessments and benchmarking, that it allows for performing a meta-analysis, and that it contributes to the evidence-based literature on the nursing work environment. However, its limitations must be acknowledged. Nurse leaders’ perceptions of a HWE were not assessed in this study, and the survey did not include qualitative items that could present comprehensive insights into the nursing work environment. In addition, the convenience sampling of the research population and the limited sample size, which may have impacted the representativeness of our sample and the generalizability of the findings. Larger sample sizes in subsequent research would be recommended to confirm the findings from the current study.

Conclusion

The study’s findings could assist hospitals in prioritizing the adoption of AACN HWE standards. Based on the findings of this study, a target benchmark of “good” was established for both the overall score and each standard, which indicates a good work environment as perceived by ICU nurses. In addition, interpersonal differences should be considered when developing improvement initiatives. Interventions could contribute to better work environments, health-care services, and patient experience. Regular assessments should be conducted periodically, and results should be used as a benchmark for future comparisons to monitor the implemented improvements and identify shortcomings.

Acknowledgments

The authors thank the nursing department supervisors who facilitated the distribution of the survey questionnaire.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Al-Bsheish M, Jarrar MT, Mustafa M, et al. ICU nurses’ safety performance related to respect for safety and management commitment: a cross-sectional study. *Contemp Nurse*. 2022;1–14. doi:10.1080/10376178.2022.2104740
2. Woodrow P. *Intensive Care Nursing: A Framework for Practice*. Routledge; 2018.

3. Rosa WE, Ferrell BR, Wiencek C. Increasing critical care nurse engagement of palliative care during the COVID-19 pandemic. *Crit Care Nurse*. 2020;40(6):e28–e36. doi:10.4037/ccn2020946
4. Sumner S. Impact of the COVID-19 pandemic on the work environment and mental health of intensive care unit nurses: reflections from the United States. *Nurs Crit Care*. 2022. doi:10.1111/nicc.12759
5. Nikbakht Nasrabadi A, Abbasi S, Mardani A, Maleki M, Vlasisavljevic Z. Experiences of intensive care unit nurses working with COVID-19 patients: a systematic review and meta-synthesis of qualitative studies. *Front Public Health*. 2022;10. doi:10.3389/fpubh.2022.1034624
6. Ulrich BT, Lavandero R, Woods D, Early S. Critical care nurse work environments 2013: a status report. *Crit Care Nurse*. 2014;34(4):64–79. doi:10.4037/ccn2014731
7. Ulrich B, Barden C, Cassidy L, Varn-Davis N. Critical care nurse work environments 2018: findings and implications. *Crit Care Nurse*. 2019;39(2):67–84. doi:10.4037/ccn2019605
8. ISO Standard. Ergonomics principles in the design of work systems; 2016.
9. American Association of Critical-Care Nurses. AACN standards for establishing and sustaining healthy work environments: a journey to excellence. *Am J Crit Care*. 2005;14(3):187–197. doi:10.4037/ajcc2005.14.3.187
10. Janíková E, Zeleníková R, Jarošová D, Plevová I, Mynaříková E. Work environment assessment instruments used in nursing. *Kontakt*. 2021;23(4):263–273. doi:10.32725/kont.2021.041
11. Institute of Medicine. *The Future of Nursing: Leading Change, Advancing Health*. Washington (DC): National Academies Press (US); 2011.
12. Efsthathiou G, Andreou C, Tsangari H, Dimitriadou M, Papastavrou E. Adaptation and validation of the cyprus version of the practice environment scale of the nursing work index: a methodological study. *BMC Res Notes*. 2018;11(1):791. doi:10.1186/s13104-018-3896-2
13. Connor JA, Zinzel SI, Porter C, et al. Interprofessional use and validation of the AACN healthy work environment assessment tool. *Am J Crit Care*. 2018;27(5):363–371. doi:10.4037/ajcc2018179
14. Institute of Medicine. *Keeping Patients Safe: Transforming the Work Environment of Nurses*. Washington (DC): National Academies Press (US); 2004.
15. Jarrar M, Al-Bsbeish M, Aldhmadi B, et al. Effect of practice environment on nurse reported quality and patient safety: the mediation role of person-centeredness. *InHealthcare*. 2021;9:1578. doi:10.3390/healthcare9111578
16. Nasirizad Moghadam K, Chehrzad MM, Reza Masouleh S, et al. Nursing physical workload and mental workload in intensive care units: are they related? *Nurs Open*. 2021;8(4):1625–1633. doi:10.1002/nop.2.785
17. Blake N, Sandoval R, Sangalang R, et al. Roadmap for improving nursing excellence using AACN's healthy work environment standards. *AACN Adv Crit Care*. 2022;33(2):208–211. doi:10.4037/aacnacc2022632
18. Purdy N, Spence Laschinger HK, Finegan J, Kerr M, Olivera F. Effects of work environments on nurse and patient outcomes. *J Nurs Manag*. 2010;18(8):901–913. doi:10.1111/j.1365-2834.2010.01172.x
19. Wei H, Sewell KA, Woody G, Rose MA. The state of the science of nurse work environments in the United States: a systematic review. *Int J Nurs Sci*. 2018;5(3):287–300. doi:10.1016/j.ijnss.2018.04.010
20. Moghadam KN, Chehrzad MM, Masouleh SR, et al. Nursing workload in intensive care units and the influence of patient and nurse characteristics. *Nurs Crit Care*. 2021;26(6):425–431. doi:10.1111/nicc.12548
21. Faul F, Erdfelder E, Lang AG, Buchner A. G*Power s3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods*. 2007;39(2):175–191. doi:10.3758/BF03193146
22. Nayback-Beebe AM, Forsythe T, Funari T, et al. Using evidence-based leadership initiatives to create a healthy nursing work environment. *Dimens Crit Care Nurs*. 2013;32(4):166–173. doi:10.1097/DCC.0b013e3182998121
23. Huddleston P, Gray J. Measuring nurse leaders' and direct care nurses' perceptions of a healthy work environment in an acute care setting, part 1: a pilot study. *J Nurs Adm*. 2016;46(7–8):373–378. doi:10.1097/NNA.0000000000000361
24. Aboshaiqah AE. Nursing work environment in Saudi Arabia. *J Nurs Manag*. 2015;23(4):510–520. doi:10.1111/jonm.12164
25. Jarrar M, Al-Bsbeish M, Dardas L, Meri A, Minai MS. Adverse events in Malaysia: associations with nurse's ethnicity and experience, hospital size, accreditation, and teaching status. *Int J Health Plan Manag*. 2020;35:104–119. doi:10.1002/hpm.2822
26. Henkin S, Chon TY, Christopherson ML, Halvorsen AJ, Worden LM, Ratelle JT. Improving nurse-physician teamwork through interprofessional bedside rounding. *J Multidiscip Healthc*. 2016;9:201–205. doi:10.2147/JMDH.S106644
27. Haykal T, Al-Dulaimi R, Sidahmed S, et al. Understanding the means of communication between nurses and resident physicians in the modern world: a community-based university hospital survey results. *J Comm Hosp Intern Med Perspect*. 2020;10(2):107–110. doi:10.1080/20009666.2020.1751515
28. Schwarzkopf D. Übertherapie – eine Belastung für Behandler und Angehörige [Nonbeneficial care-a burden for clinicians and relatives]. *Med Klin Intensivmed Notfmed*. 2019;114(3):222–228. German. doi:10.1007/s00063-019-0531-5
29. Bhutani J, Bhutani S, Balhara YP, Kalra S. Compassion fatigue and burnout amongst clinicians: a medical exploratory study. *Indian J Psychol Med*. 2012;34(4):332–337. doi:10.4103/0253-7176.108206
30. Al-Mugheed K, Bayraktar N, Al-Bsbeish M, et al. Patient safety attitudes among doctors and nurses: associations with workload, adverse events, experience. *Healthcare*. 2022;10(4):1. doi:10.3390/healthcare10040631
31. Grissinger M. Disrespectful behavior in health care: its impact, why it arises and persists, and how to address it-part 2. *PT*. 2017;42(2):74–77.
32. Rosenstein AH, O'Daniel M. A survey of the impact of disruptive behaviors and communication defects on patient safety. *Jt Comm J Qual Patient Saf*. 2008;34(8):464–471. doi:10.1016/s1553-7250(08)34058-6
33. McNamara SA. Incivility in nursing: unsafe nurse, unsafe patients. *AORN J*. 2012;95(4):535–540. doi:10.1016/j.aorn.2012.01.020
34. Boev C, Xia Y. Nurse-physician collaboration and hospital-acquired infections in critical care. *Crit Care Nurse*. 2015;35(2):66–72. doi:10.4037/ccn2015809
35. Maxfield D, Grenny J, McMillan R, Patterson K, Switzler A. *Silence Kills: The Seven Crucial Conversations in Healthcare*. Aliso Viejo, CA: American Association of Critical-Care Nurses (AACN) and Vital Smarts; 2005.
36. Hunsaker S, Chen HC, Maughan D, Heaston S. Factors that influence the development of compassion fatigue, burnout, and compassion satisfaction in emergency department nurses. *J Nurs Scholarsh*. 2015;47(2):186–194. doi:10.1111/jnu.12122
37. Sacco TL, Czurzynski SM, Harvey ME, Ingersoll GL. Compassion satisfaction and compassion fatigue among critical care nurses. *Crit Care Nurse*. 2015;35(4):32–43. doi:10.4037/ccn2015392

38. Robertson-Malt S, Chapman Y. Finding the right direction: the importance of open communication in a governance model of nurse management. *Contemp Nurse*. 2008;29(1):60–66.
39. Al-Bsheish M, Mustafa M, Ismail M, Jarrar M, Meri A, Dauwed M. Perceived management commitment and psychological empowerment: a study of intensive care unit nurses' safety. *Saf Sci*. 2019;118:632–640. doi:10.1016/j.ssci.2019.05.055
40. Shin S, Park JH, Bae SH. Nurse staffing and nurse outcomes: a systematic review and meta-analysis. *Nurs Outlook*. 2018;66(3):273–282. doi:10.1016/j.outlook.2017.12.002
41. Jarrar M, Rahman H, Minai MS, Abumadini M, Larbi M. The function of patient-centered care in mitigating the effect of nursing shortage on the outcomes of care. *Int J Health Plann Manag*. 2018;33:1.
42. Jarrar M, Rahman H, Sebiany A, Abumadini M, Masnawaty H, Vallabadoss CA. Nursing duty hours' length and the perceived outcomes of care. *Glob J Health Sci*. 2018;10:1.
43. McClelland LE, Vogus TJ. Compassion practices and HCAHPS: does rewarding and supporting workplace compassion influence patient perceptions? *Health Serv Res*. 2014;49(5):1670–1683. doi:10.1111/1475-6773.12186
44. Lefton C. Strengthening the workforce through meaningful recognition. *Nurs Econ*. 2012;30(6):331–338.
45. Ray SL, Wong C, White D, Heaslip KJT. Compassion satisfaction, compassion fatigue, work life conditions, and burnout among frontline mental health care professionals. *Traumatology*. 2013;19(4):255–267. doi:10.1177/1534765612471144

Risk Management and Healthcare Policy

Dovepress

Publish your work in this journal

Risk Management and Healthcare Policy is an international, peer-reviewed, open access journal focusing on all aspects of public health, policy, and preventative measures to promote good health and improve morbidity and mortality in the population. The journal welcomes submitted papers covering original research, basic science, clinical & epidemiological studies, reviews and evaluations, guidelines, expert opinion and commentary, case reports and extended reports. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/risk-management-and-healthcare-policy-journal>