

# Inappropriate Analysis of a Cluster Randomized Controlled Trial Due to Not Accounting for Nesting and Clustering [Response to Letter]

Daphne Sze Ki Cheung<sup>1</sup>, Paul Hong Lee<sup>2</sup>, Claudia Kam Yuk Lai<sup>1</sup>

<sup>1</sup>School of Nursing, The Hong Kong Polytechnic University, Hong Kong; <sup>2</sup>Southampton Clinical Trials Unit, University of Southampton, Southampton, UK

Correspondence: Daphne Sze Ki Cheung, Email [daphne.cheung@polyu.edu.hk](mailto:daphne.cheung@polyu.edu.hk)

## Dear editor

We appreciate the interest of the readers regarding the analysis of our study entitled “A home-based dyadic music-with-movement intervention for people with dementia and caregivers: a hybrid type 2 cluster-randomized effectiveness-implementation design”.<sup>1</sup> We are writing to respond to the criticism of the validity of the conclusion of our paper. Here, we report the results of the additional analysis to address the potential clustering effects of the intervention on the outcomes as suggested by Hefner et al.<sup>2</sup>

Our initial published analysis has taken reference to similar studies and relevant guidelines. The objective of our study was to explore the population-average effect (i.e., the impact of the intervention on the participants). Thus, using the generalized estimating equation (GEE) was an appropriate analysis for outcomes variables across the two time points (baseline and follow-up), regardless of the normality of distribution.<sup>3</sup> Before conducting the GEE, we confirmed no cluster effect on the major primary outcomes at baseline by using the GLM univariate analysis. The primary outcomes of interest are the anxiety and depressive symptoms of people with dementia, and the perceived stress of caregivers, which showed no significant variation among centres (see Table 1). Hence, we performed the GEE analysis without considering centre as an independent variable. Various teams of authors adopted a similar analytical approach too.<sup>4,5</sup>

However, we notice that a sub-scale outcome (Cornell Scale for Depression in Dementia – Ideational subscale) and a secondary outcome (Quality of the Caregiver–Care Recipient Relationship) are significantly different across centres, using an uncorrected *p*-value cut-off of 0.05, when we revisit the data. Therefore, after discussing with statisticians, we performed a sensitivity analysis by adjusting centres. Similar to the initial published analysis results,<sup>1</sup> the GEE analysis adjusted to affiliated centre results showed that participants with dementia in the intervention group had improvement in anxiety status and depressive symptoms; and caregivers in the intervention group also had lowered level of perceived stress as compared to the control group (Table 2).

Concerning the data sharing request, we have indicated in the Data Sharing Statement section that the datasets are not publicly available due to ethical restrictions. The journal’s data-sharing policy allows the authors to share or not to share the data consistent with the terms of consent signed by study participants. We did not obtain consent from our participants to share the data beyond the research team, so unfortunately, Ms Hefner’s request of the datasets could not be fulfilled. Although we were not asked to provide details about our methods in the email by Ms Hefner in October 2022, as stated in the published Letter to Editor, we trust that this response letter has addressed their query.

There are different ways of handling statistics as long as they are justifiable and help researchers to understand the data, that is why we had never received similar doubts from the reviewers or journal editors. As the late George Box (1976), a British statistician regarded by many as one of the great statistical minds of the 20th century, mentioned, “All models are wrong, but some are useful”.<sup>6</sup> We are thankful to the journal for letting us publish the additional data analysis results to clarify the concern being raised.

**Table 1** Comparing Outcomes at Baseline across Centres

		Intervention (n = 55)				Control (n = 45)			p-value
		Centre 1	Centre 2	Centre 3	Centre 6	Centre 4	Centre 5	Centre 7	
		Mean ± SD or %				Mean ± SD or Count (%)			
RAID total score	Baseline	12.5 ± 5.8	10.3 ± 6.6	11.2 ± 11.7	12.1 ± 8.1	17.3 ± 11.3	10.3 ± 5.6	11.0 ± 9.4	0.469
	FU	8.5 ± 7.9	7.5 ± 8.5	10.6 ± 11.8	7.5 ± 6.1	15.5 ± 10.9	9.3 ± 5.2	11.0 ± 6.2	
Being anxious	Baseline	70%	41.7%	35.7%	57.9%	66.7%	38.5%	40.0%	0.386
	FU	25%	18.2%	33.3%	31.3%	70%	55.6%	55.6%	
CSDD total score	Baseline	5.5 ± 2.3	5.9 ± 4.6	5.4 ± 5.5	9.0 ± 5.2	10.3 ± 7.0	6.6 ± 3.9	8.1 ± 4.3	0.073
	FU	6.5 ± 5.5	2.6 ± 2.4	7.0 ± 6.6	5.3 ± 4.6	12.1 ± 7.0	7.8 ± 3.9	8.4 ± 4.4	
Being depressive <sup>###</sup>	Baseline	0%	16.7%	7.1%	36.8%	41.7%	15.4%	20%	0.100
	FU	25%	0%	25.0%	18.8%	50.0%	11.1%	27.8%	
CSDD_Mood	Baseline	0.5 ± 0.7	1.3 ± 1.7	1.6 ± 2.1	2.1 ± 2.2	2.7 ± 2.0	1.6 ± 1.2	2.0 ± 1.7	0.137
	FU	1.3 ± 2.5	0.6 ± 0.8	1.9 ± 1.8	1.3 ± 1.7	3.6 ± 1.8	2.2 ± 1.7	1.8 ± 1.5	
CSDD_Behavioural	Baseline	1.8 ± 1.4	1.5 ± 1.0	2.1 ± 2.1	2.2 ± 1.6	2.3 ± 1.8	1.4 ± 1.0	2.7 ± 1.5	0.284
	FU	2.0 ± 1.8	0.9 ± 0.8	2.2 ± 2.5	0.7 ± 0.8	2.5 ± 1.6	2.0 ± 1.2	2.2 ± 1.4	
CSDD_Physical	Baseline	1.6 ± 1.4	1.1 ± 1.1	0.4 ± 0.7	1.1 ± 1.1	1.2 ± 1.0	0.9 ± 1.0	1.2 ± 1.3	0.282
	FU	0.5 ± 0.6	0.6 ± 0.8	1.1 ± 1.1	0.5 ± 0.9	1.4 ± 0.8	0.9 ± 1.1	1.4 ± 1.2	
CSDD_Cyclic	Baseline	1.6 ± 1.2	1.3 ± 1.4	0.5 ± 1.2	2.2 ± 2.1	1.9 ± 2.0	1.9 ± 2.2	1.2 ± 0.9	0.097
	FU	2.0 ± 2.8	0.2 ± 0.4	0.1 ± 0.4	1.8 ± 1.3	1.7 ± 1.6	1.6± 1.5	1.6 ± 1.5	
CSDD_Ideational	Baseline	0.0 ± 0.0	0.8 ± 1.4	0.7 ± 1.0	1.5 ± 1.8	2.3 ± 2.5	0.9 ± 1.2	1.1 ± 1.6	0.026
	FU	0.8 ± 1.0	0.5 ± 0.8	1.1 ± 2.3	1.0 ± 1.5	2.9 ± 3.2	1.1 ± 1.5	1.4 ± 1.3	
PSS total score	Baseline	15.2 ± 4.0	15.8 ± 3.9	14.4 ± 8.1	18.7 ± 4.6	19.0 ± 6.4	15.9 ± 6.3	19.2 ± 5.0	0.097
	FU	17.5 ± 3.3	12.6 ± 7.5	15.4 ± 7.7	14.8 ± 5.3	19.9 ± 7.1	16.5 ± 5.0	19.9 ± 4.3	
PAC total score	Baseline	34.3 ± 8.3	35.5 ± 6.6	36.9 ± 6.8	34.8 ± 8.3	28.6 ± 8.7	35.5 ± 4.9	31.7 ± 8.0	0.090
	FU	34.7 ± 7.8	34.0 ± 5.1	32.7 ± 9.7	36.5 ± 8.3	32.6 ± 16.3	34.2 ± 8.0	33.2 ± 8.2	
QCCRR total score	Baseline	11.2 ± 2.5	13.0 ± 2.7	12.3 ± 2.6	11.5 ± 1.6	10.3 ± 2.4	11.1 ± 1.8	10.3 ± 2.0	0.015
	FU	12.0 ± 1.4	12.5 ± 3.2	11.8 ± 3.4	12.1 ± 2.3	10.6 ± 1.71	11.7 ± 2.4	10.7 ± 2.8	

Notes: <sup>#</sup>RAID\_Total > 10 indicates anxious; <sup>###</sup>CSDD\_Total > 10 indicates depressive.

Abbreviations: FU, Follow-up; RAID, Rating Anxiety in Dementia; CSDD, Cornell Scale for Depression in Dementia; PSS, Perceived Stress Scale; PAC, Positive Aspect of Caregiving; QCCRR, Quality of the Caregiver–Care Recipient Relationship.

**Table 2** Group  $\times$  Time GEE Analysis Results Adjusted the Clustering Effects

Outcomes	B (SE)	95% Wald CI [Lower, Upper]	p-value
RAID total score	−2.78 (1.74)	[−6.20, 0.64]	0.111
Anxious or not	−1.60 (0.52)	[−2.60, −0.59]	0.002*
CSDD total score	−3.10 (1.06)	[−5.19, −1.02]	0.004*

(Continued)

Table 2 (Continued).

Outcomes	B (SE)	95% Wald CI [Lower, Upper]	p-value
Depressive or not	-0.56 (0.64)	[-1.82, 0.70]	0.382
CSDD_Mood	-0.72 (0.35)	[-1.41, -0.03]	0.040*
CSDD_Behavioural	-0.83 (0.36)	[-1.53, -0.12]	0.021*
CSDD_Physical	-0.50 (0.29)	[-1.06, 0.06]	0.079
CSDD_Cyclic	-0.50 (0.39)	[-1.26, 0.25]	0.192
CSDD_Ideational	-0.55 (0.36)	[-1.25, 0.15]	0.124
PSS total score	-2.45 (1.20)	[-4.81, -0.10]	0.041*
PAC total score	-2.10 (2.17)	[-6.34, 2.15]	0.332
QCCRR total score	-0.39 (0.53)	[-1.43, 0.65]	0.458

Note: \*Statistically significant.

## Funding

The primary study was funded by Ho Cheung Shuk Yuen Charitable Foundation and this additional analysis received no funding support.

## Disclosure

The authors report no conflicts of interest in this communication.

## References

- Cheung DSK, Ho LYW, Chan LCK, Kwok RKH, Lai CKY. A home-based dyadic music-with-movement intervention for people with dementia and caregivers: a hybrid type 2 cluster-randomized effectiveness-implementation design. *Clin Interv Aging*. 2022;11(17):1199–1216. doi:10.2147/CIA.S370661
- Hefner M, Jamshidi-Naeini Y, Vorland C, Golzarri-Arroyo L, Shaw B, Allison D. Inappropriate analysis of a cluster randomized controlled trial due to not accounting for nesting and clustering: comment on “A home-based dyadic music-with-movement intervention for people with dementia and caregivers: a hybrid type 2 cluster randomized effectiveness-implementation design”. *Clin Interv Aging*. 2022;2022:1.
- Diggle PJ, Heagerty P, Liang KY, Zeger SL. *Analysis of Longitudinal Data*. 2nd ed. Oxford, UK: Oxford University Press; 2002.
- Lin YC, Wang CJ, Chang YF, Wang JJ. Effects of the biopsychosocial functional activity program on cognitive function for community older adults with mild cognitive impairment: a cluster-randomized controlled trial. *Nurs Health Sci*. 2020;22(4):1065–1075. doi:10.1111/nhs.12772
- Chen SM, Lin HS, Atherton JJ, MacIsaac RJ, Wu CJ. Effect of a mindfulness programme for long-term care residents with type 2 diabetes: a cluster randomised controlled trial measuring outcomes of glycaemic control, relocation stress and depression. *Int J Older People Nurs*. 2020;15(3):e12312. doi:10.1111/opn.12312
- Box GE. Science and statistics. *J Am Stat Assoc*. 1976;71(356):791–799. doi:10.1080/01621459.1976.10480949

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Clinical Interventions in Aging ‘letters to the editor’ section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Clinical Interventions in Aging editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Clinical Interventions in Aging

Dovepress

Publish your work in this journal

Clinical Interventions in Aging is an international, peer-reviewed journal focusing on evidence-based reports on the value or lack thereof of treatments intended to prevent or delay the onset of maladaptive correlates of aging in human beings. This journal is indexed on PubMed Central, MedLine, CAS, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-interventions-in-aging-journal>